## the Handy Bt Trait Table

for U.S. Corn Production

**Updated** January 2017

Posted at www.msuent.com

For questions, complaints, or corrections: Chris DiFonzo, Michigan State University, difonzo@msu.edu Contributors: Pat Porter, Texas A&M University & Kelley Tilmon, The Ohio State University

Most corn hybrids planted in the U.S. contain one or more transgenic traits for weed or insect management. These traits can increase flexibility and profitability for producers, but sometimes cause confusion about their spectrum of control or refuge requirements. The Handy Bt Trait Table provides a helpful list of trait names (below) and details of trait packages (next page) to make it easier to read company seed guides, sales materials, and bag tags. Note that there are two versions of the table (north/Midwest vs. south/cotton belt) which differ only in refuge percentages.

## Important clarifications or changes to the Trait Table for 2017

- ✓ An insect is listed in the CONTROL SPECTRUM column if seed providers claim protection or efficacy for a given Bt package; insect species which are 'suppressed' are no longer listed. Actual field-level performance of hybrids on lepidopteran and rootworm larvae may differ if there are local or regional insect populations which are less susceptible or resistant to Bt proteins.
- √ To address local or regional performance issues, a new column ('May be Ineffective On') was added to highlight insect x Bt combinations with documented field-failures, confirmed resistance, or cross-resistance. An insect is listed in this column only if ALL of the Bt proteins which should control it in a product are 'ineffective' somewhere in the US or Canada. Ineffective ratings are based on published lab assays &/or field research from field corn, sweet corn, and cotton. University extension specialists or local educators can assist in determining if you are in an area where reduced effectiveness was reported. On a broader scale, this column is intended to alert growers and consultants to potential management problems, influence seed selection, and encourage field scouting.
- ✓ The refuge column was simplified to include only the % and an indication if the refuge is in the bag.

Field corn 'events' (transformations of one or more genes) and their Trade Names

Trade name for trait	Event	Protein(s) expressed	Insect Target or Herbicide Activity	
Agrisure CB/LL	Bt11	Cry1Ab + <i>PAT</i>	corn borer + glufosinate tolerance	
Agrisure Duracade	5307	eCry3.1Ab	rootworm	
Agrisure GT	GA21	EPSPS	glyphosate tolerance	
Agrisure RW	MIR604	mCry3A	rootworm	
Agrisure Viptera	MIR162	Vip3A	broad lep control (but not corn borer)	
Herculex 1 or CB	TC1507	Cry1Fa2 + <i>PAT</i>	corn borer + glufosinate tolerance	
Herculex RW	DAS-59122-7	Cry34Ab1/Cry35Ab1 + <i>PAT</i>	rootworm + glufosinate tolerance	
Roundup Ready 2	NK603	EPSPS	glyphosate tolerance	
Yieldgard Corn Borer	MON810	Cry1Ab	corn borer	
Yieldgard Rootworm	MON863	Cry3Bb1	rootworm	
Yieldgard VT Pro	MON89034	Cry1A.105 + Cry2Ab2	broader lep control	
Yieldgard VT Rootworm RR	MON88017	Cry3Bb1 + EPSPS	rootworm + glyphosate tolerance	

## Abbreviations used in the Trait Table

Insect targets BCW black cutworm CEW corn earworm SB stalk borer ECB European corn borer SWCB southwestern corn borer FAW fall armyworm TAW true armyworm RW corn rootworm WBC western bean cutworm

Herbicide activity

LL

dicamba tolerant

GT alyphosate tolerant

Liberty Link - glufosinate-tolerant RR2 Roundup Ready 2, glyphosate-tolerant

TDAIT FARALLY	1	CONTROL SPECTRUM		Marria Institu		
TRAIT FAMILY		CONTROL SPECTRUM	May be locally	l la mbiaida	Refuge -	
Considia Dua durat		Marketed for control of	or regionally	Herbicide	Midwest/	
Specific Product AGRISURE	Bt protein(s)	above-ground	-in soii	ineffective on:	tolerance	North
Agrisure 3010, 3010A	Cry1Ab	ECB SWCB	ı		GT LL	20%
Agrisure 3000GT, 3011A	Cry1Ab	ECB SWCB	I RW	RW	GT LL	20%
Agrisure 3000GT, 3011A	mCry3A	ECB 3WCB	i Kvv	L VV	GI LL	20%
Agrisure Viptera 3110	Cry1Ab Vip3A	BCW CEW ECB FAW	l		GT LL	20%
Agrisure viptera 3110	Crylab Vipsa		i		GI LL	20%
Agrisure Viptera 3111	Cry1Ab Vip3A	SB SWCB TAW WBC BCW CEW ECB FAW	I RW	RW	GT LL	20%
Agrisure viptera 3111	mCry3A		I	I K V V	GI LL	20%
Agrisure	Cry1Ab Cry1F	SB SWCB TAW WBC BCW ECB FAW	l RW	FAMAL MADC	GT	FO/ in hor
_	1 '		i Kvv	FAW, WBC	GI	5% in bag
3122 E-Z Refuge	mCry3A Cry34/35Ab1	SB SWCB WBC BCW CEW ECB FAW	<u> </u>	RW	GT	FO/ in hor
Agrisure Viptera 3220 E-Z Refuge	Cry1Ab Cry1F Vip3A	SB SWCB TAW WBC	·		GI	5% in bag
Agrisure Duracade	Cry1Ab Cry1F	BCW ECB FAW	RW	FAW, WBC	GT	5% in bag
-	1 ' ·		I LAA	RW	Gi	5% III Dag
5122 E-Z Refuge Agrisure Duracade	mCry3A eCry3.1Ab Cry1Ab Cry1F Vip3A	SB SWCB WBC  BCW CEW ECB FAW	RW	RW	GT	5% in bag
5222 E-Z Refuge	1 ' ' '	SB SWCB TAW WBC	I LAA	L VV	Gi	5% III Dag
HERCULEX	mCry3A eCry3.1Ab	3B 3WCB IAW WBC	-			
Herculex 1 (HX1)	Cry1F	BCW ECB FAW	l I	FAW, SWCB, WBC	LL	20%
Herculex 1 (HX1)	CIVIF	SB SWCB WBC		FAW, SWCB, WBC	LL	20%
Herculex RW (HXRW)	Cry34/35Ab1		l RW	RW	RR2	20%
Herculex XTRA (HXX)	Cry1F	BCW ECB FAW	i RW	FAW, SWCB, WBC		20%
Herculex ATRA (HAA)	Cry34/35Ab1	SB SWCB WBC	I NVV	RW	(most)	20%
OPTIMUM	C1 y 34 / 3 3 A D 1	3B 3WCB WBC		I N V V		
Intrasect (YHR)	Cry1Ab Cry1F	BCW ECB FAW	<u>.</u>	FAW, WBC	LL RR2	5%
intrasect (TTIIV)	CIVIAD CIVII	SB SWCB WBC	I	TAW, WBC	LL IXIXZ	370
AcreMax (AM)	Cry1Ab Cry1F	BCW ECB FAW	<del></del>	FAW, WBC	LL RR2	5% in bag
ACICIVIAX (AIVI)	CIVIAD CIVII	SB SWCB WBC	 	TAW, WBC	LL IXIXZ	370 III Dag
Leptra (VYHR) <sup>a</sup>	Cry1Ab Cry1F Vip3A	BCW CEW ECB FAW	· 		LL RR2	a5% ⅓ mile
AcreMax Leptra (AML) <sup>b</sup>	CIVIAD CIVIF VIDSA	SB SWCB TAW WBC	 		LL KKZ	b5% in bag
AcreMax RW (AMRW)	Cry34/35Ab1		RW	RW	LL RR2	10% in bag
AcreMax1 (AM1)	Cry1F	BCW ECB FAW	RW	FAW, SWCB, WBC	LL RR2	10% in bag
THE CIVILAT (THE I)	Cry34/35Ab1	SB SWCB WBC	! '`'	RW	LL IIIL	20% ECB
TRIsect (CHR)	Cry1F	BCW ECB FAW	RW	FAW, SWCB, WBC	LL RR2	20% ECB
Thiseet (erm)	mCry3A	SB SWCB WBC	I ''''	RW	LL IIIL	2070
Intrasect TRIsect (CYHR) <sup>a</sup>	Cry1Ab Cry1F	BCW ECB FAW	RW	FAW, WBC	LL RR2	a20%
AcreMax TRIsect (AMT) <sup>b</sup>	mCry3A	SB SWCB WBC	'\'V	RW	LL IVIVE	<sup>b</sup> 10% in bag
Intrasect Xtra (YXR) <sup>a</sup>	Cry1Ab Cry1F	BCW ECB FAW	RW	FAW, WBC	LL RR2	<sup>a</sup> 20%
AcreMax Xtra (AMX) <sup>b</sup>	Cry34/35Ab1	SB SWCB WBC	 	RW		b10% in bag
Intrasect Xtreme (CYXR) <sup>a</sup>	Cry1Ab Cry1F		RW	FAW, WBC	LL RR2	a5%
AcreMax XTreme (AMXT) <sup>b</sup>	mCry3A Cry34/35Ab1	SB SWCB WBC	i	RW	LL IIIL	<sup>b</sup> 5% in bag
YIELDGARD or GENUITY	merysit erystyssitsi	35 3WC5 WBC		11.00		370 III Bug
YieldGard CB (YGCB)	Cry1Ab	ECB SWCB	<u> </u>		RR2	20%
YieldGard VT Rootworm	Cry3Bb1		ı RW	RW	RR2	20%
YieldGard VT Triple	Cry1Ab	ECB SWCB	RW	RW	RR2	20%
Ticiadara VI Tipic	Cry3Bb1	Leb Sweb	 		MAZ	2070
Genuity VT Double PRO <sup>a</sup>	Cry1A.105 Cry2Ab2	CEW ECB FAW SB SWCB		CEW	RR2	a5%
or RIB complete <sup>b</sup>	,		i			<sup>b</sup> 5% in bag
Genuity VT Triple PRO <sup>a</sup>	Cry1A.105 Cry2Ab2	CEW ECB FAW SB SWCB	RW	CEW	RR2	<sup>a</sup> 20%
or RIB complete <sup>b</sup>	Cry3Bb1		i	RW		<sup>b</sup> 10% in bag
Genuity SmartStax <sup>a</sup>	Cry1A.105 Cry2Ab2 Cry1F	BCW CEW ECB FAW	RW	CEW, WBC	LL RR2	a5%
or RIB Complete <sup>b</sup>	Cry3Bb1 Cry34/35Ab1	SB SWCB WBC	<b></b>	RW		<sup>b</sup> 5% in bag
OTHER	. ,					
Powercore <sup>a</sup>	Cry1A.105 Cry2Ab2 Cry1F	BCW CEW ECB FAW	·	CEW, WBC	LL RR2	<sup>a</sup> 5%
Powercore Refuge Adv.b	, , , , , , , , , , , , , , , , , , , ,	SB SWCB WBC	<u>'</u>			<sup>b</sup> 5% in bag
Smartstax <sup>a</sup>	Cry1A.105 Cry2Ab2 Cry1F	BCW CEW ECB FAW	RW	CEW, WBC	LL RR2	a5%
Smartstax Refuge Adv.b	Cry3Bb1 Cry34/35Ab1	SB SWCB WBC	, I	RW		<sup>b</sup> 5% in bag
	, ,,				L	