

Archer, S, TW Boutton, KA Hibbard. 2001. Trees in grasslands: biogeochemical consequences of woody plant expansion, pp. 115-137. In: *Global Biogeochemical Cycles in the Climate System* (E-D Schulze, SP Harrison, M Heimann, EA Holland, J Lloyd, IC Prentice, D Schimel, eds.). Academic Press, San Diego.

Trees in Grasslands: Biogeochemical Consequences of Woody Plant Expansion

Steve Archer,
Thomas W. Boutton,
and K. A. Hibbard
Texas A&M University
College Station,
Texas

1. Introduction	115
2. Woody Plant Encroachment in Grasslands and Savannas	116
3. The La Copita Case Study	118
4. Degradation: Ecological versus Socioeconomic	127
5. Implications for Ecosystem and Natural Resources Management	128
6. Summary	129
7. References	130

Key Words: carbon, disturbance, grazing, fire, global change, hydrocarbons, land cover, modeling, nitrogen, NO_x, savanna, sequestration, soil respiration, succession, tree-grass interactions, vegetation dynamics, woody plant encroachment

1. Introduction

The term "savanna" typically denotes plant communities or landscapes having a continuous grass layer with scattered woody plants. Although savannas are not the only vegetation type where contrasting plant life forms codominate, they are one of the most striking, geographically extensive (ca. 20% of global land surface; Scholes and Hall, 1996) and socioeconomically important examples in tropical (Tothill and Mott, 1985; Young and Solbrig, 1993) and temperate (Burgess, 1995; McPherson, 1997; Anderson *et al.*, 1999) regions. Tropical savannas cover about 1600 million ha of the terrestrial surface (Scholes and Hall, 1996), including more than half the area of Africa and Australia, 45% of South America, and 10% of India and southeastern Asia (Werner, 1991). Temperate savannas in North America occupy an estimated 50 million ha (McPherson, 1997). More importantly, savannas contain a large and rapidly growing proportion of the world's human population and a majority of its rangelands and domesticated animals. As such, they have received substantial and ever-increasing anthropogenic land-use pressure.

Many savannas are dynamic ecotones between woody plant (shrub-steppe, desert scrub, woodland or forest) and grassland formations. Savannas vary substantially with respect to the stature

(shrub vs tree), canopy cover (e.g., 5–80%), functional form (evergreen vs deciduous; broad-leaved vs needle-leaved; shallow vs deeply rooted), and spatial arrangement (random, regular, or clumped) of the woody elements that compose them. Similarly, the grass layer may consist of short versus tall-statured species, bunch versus rhizomatous growth forms, and C₃, C₄, or mixed C₃/C₄ photosynthetic pathway assemblages. This variation in structural/functional characteristics reflects a rich array of interactions between climate (especially the amount and seasonality of rainfall), soils (notably, depth and texture), and disturbance (particularly grazing, browsing, and fire), as shown in Figure 1 (Walker, 1987; Backéus, 1992).

Much of the literature on savanna ecology has been devoted to describing and classifying vegetation structure. Static classification schemes minimize the importance of temporal change and divert attention from functional processes that might explain dynamic spatiotemporal variation. Grasses and woody plants may coexist in a dynamic equilibrium when climatic, edaphic, and disturbance factors interact temporally such that neither life form can exclude the other. However, a directional change in one or more of these primary controlling factors may shift the balance in favor of one life form over the other and move the system toward either grassland or shrubland/woodland. The probability, rate, and extent of such a shift may depend on local topoeconomic factors and the life-history traits and autecology of the growth forms or species involved.

Human population growth and widespread Anglo-European settlement during the 18th and 19th centuries have influenced the balance of grass-woody plant interactions worldwide. For example, extensive clearing of trees for fuel, lumber, and cropland has fragmented forests and produced anthropogenic or degraded savannas (Gadgil and Meher-Homji, 1985; Sinclair and Fryxell, 1985; Cline-Cole *et al.*, 1990; Schüle, 1990; Young and Solbrig,

¹ Current Address: Climate Change Research Center, GAIM Task Force, Institute for the Study of Earth, Oceans, and Space (EOS), Morse Hall, 39 College Road, University of New Hampshire, Durham, NH 03824-3525




FIGURE 1 Numerous factors interact to affect the abundance of grasses and woody vegetation in drylands (from Scholes and Walker, 1993). The balance between trees and grasses (innermost level) is affected by determinants of structure and function (water, nutrients, fire, and herbivory). The outermost level contains the factors that give the determinants their characteristics. Over the past century, human influences have shifted the balance to favor woody plants through selective utilization of grasses by livestock maintained at high concentrations, elimination of browsers, and fire suppression (see Archer, 1994).

1993; Mearns, 1995). Following forest clearing, pyrophytic grasses may establish and restrict woody colonization by accelerating fire cycles and maintaining low-fertility soils (Hopkins, 1983; Mueller-Dombois and Goldammer, 1990; D'Antonio and Vitousek, 1992). In the Brazilian Cerrado, rates of agricultural expansion and clearing of savanna and woodland trees rival those reported for Amazon rain forest (Klink *et al.*, 1993). In other areas, fire suppression, eradication of indigenous savanna browsers, and the introduction of grazing livestock and exotic trees and shrubs have caused a progressive increase in woody plant density, known as bush or brush encroachment (Adamoli *et al.*, 1990; Archer, 1994; Gardener *et al.*, 1990; Miller and Wigand, 1994; Noble, 1997). As a result, areas that were once forest may become savanna-like, while areas that were once grassland or open savanna may progress toward a shrubland or woodland physiognomy. The biogeochemical consequences of this latter phenomenon are the focus of this chapter.

2. Woody Plant Encroachment in Grasslands and Savannas

Woody plant encroachment has been widespread in grassland and savanna ecosystems of North and South America, Australia, Africa, and southeast Asia over the past century (Table 1). This encroachment, typically by unpalatable trees and shrubs, has gone to completion on some landscapes and is in progress on others. It jeopardizes grassland biodiversity and threatens the sustainability of pastoral, subsistence, and commercial livestock grazing (Rappole *et al.*, 1986; Noble, 1997). As such, it may adversely impact ~20% of the world's population (Turner *et al.*, 1990). The proximate causes of displacement of perennial grasses by woody plants are subjects of debate. Land-use practices such as heavy grazing and reductions in fire frequency have often been implicated. However, climate change, historic atmospheric CO₂ enrichment, and exotic species introductions are potentially important contributing factors (Idso, 1995; Archer *et al.*, 1995; Polley, 1997; Polley *et al.*, 1997). Current trends in atmospheric CO₂ enrichment may exacerbate the shifts from grass to woody plant domination, especially where the invasive trees/shrubs are capable of symbiotic N₂ fixation. Expansion of woody plants into grasslands may also be favored by recent increases in atmospheric N deposition (Köchy, 1999). In addition to influencing vegetation composition, changes in each of these factors would have the potential to alter the storage and dynamics of C and N in savanna ecosystems. The net outcome of such interactions over the recent past is poorly understood and has not been well documented.

Although woody plant encroachment has long been a concern of land managers in grassland and savanna regions (e.g., Fisher, 1950, 1977), research on this problem has been primarily "applied" and focused on the effects of woody plants on grass production and the development of chemical or mechanical methods to reduce the abundance of established trees and shrubs. Despite the long-standing recognition of woody plant encroachment as a worldwide dryland management problem, little is known of the rates and dynamics of the phenomenon or its impact on fundamental ecological processes related to energy flow, nutrient cycling, and biodiversity. Grassland/savanna systems account for 30–35% of global terrestrial net primary production (Field *et al.*, 1998). Hence, when woody species increase in abundance and transform shrublands into woodlands, grasslands into savannas, or savannas into shrublands and woodlands, the potential to alter C and N sequestration and cycling at regional and global scales may be significant. Consequently, this type of land cover change has the potential to contribute significantly to the terrestrial global carbon sink (cf. Ciais *et al.*, 1995). Savanna landforms may have a larger impact on the global carbon cycle than previously appreciated (Hall *et al.*, 1995; Ojima *et al.*, 1993; Scholes and Hall, 1996; Scholes and Bailey, 1996; Scholes and van der Merwe, 1996). Indeed, recent assessments suggest that savanna ecosystems have among the highest potential C gain and loss rates of the world's biomes (ORNL, 1998). In addition, emissions of radiatively active

TABLE 1 Survey of Studies Describing or Quantifying Woody Plant Encroachment into Grassland, Tree/Shrub Proliferation in Savannas, and Tree Encroachment into Shrubland.

Arizona	Kansas	South Dakota	Reichard and Hamilton, 1997	Burrows <i>et al.</i> , 1998
Arnold, 1950	Abrams, 1986	Bock and Bock, 1984	Rogers, 1982	Burrows <i>et al.</i> , 1990
Bahre, 1991	Bragg and Hulbert, 1976	Progulskie, 1974	Robinson, 1965	Cook <i>et al.</i> , 1996
Bahre and Shelton, 1993	Briggs and Gibson, 1992	Tieszen and Archer, 1990	Tieszen and Archer, 1990	Cunningham and Walker, 1973
Brown, 1950	Knapp and Seastedt, 1986	Tieszen and Pfau, 1995	Wall, 1999	Gardiner and Gardiner, 1996
Brown <i>et al.</i> , 1997	Knight <i>et al.</i> , 1994	Texas	West, 1988	Grice, 1996
Cooper, 1960	Loehle <i>et al.</i> , 1996	Ansley <i>et al.</i> , 1995	Young <i>et al.</i> , 1979	Grice, 1997
Covington and Moore, 1994	Owensby <i>et al.</i> , 1973	Archer <i>et al.</i> , 1988	AFRICA	Harrington <i>et al.</i> , 1979
Glendenning, 1952	Grimm, 1983	Archer, 1989	Acocks, 1964	Harrington and Hodgkinson, 1986
Humphrey and Mehrhoff, 1958	Johnston <i>et al.</i> , 1996	Boutton <i>et al.</i> , 1998	Ambrose and Sikes, 1991	Hodgkin, 1984
Hastings and Turner, 1965	Montana	Bogusch, 1952	Ben-Shaher, 1991	Lonsdale, 1993
Johnsen, 1962	Arno and Gruell, 1986	Bray, 1901	Bews, 1917	Noble, 1997
Kenney <i>et al.</i> , 1986	Arno <i>et al.</i> , 1995	Bruce <i>et al.</i> , 1995	Bond <i>et al.</i> , 1994	Panetta and McKee, 1997
Martin, 1975	Nebraska	Ellis and Schuster, 1968	Friedel, 1987	CANADA
Martin and Turner, 1977	Johnson, 1994	Foster, 1917	Grossman and Gandar, 1989	Archibald and Wilson, 1980
McClaran and McPherson, 1995	Steinauer and Bragg, 1987	Inglis, 1964	Hochberg <i>et al.</i> , 1994	Brown, 1994
McPherson <i>et al.</i> , 1993	Steuter <i>et al.</i> , 1990	Johnston, 1963	Holmes and Cowling, 1997	Köchy, 1999
Miller, 1921	New Mexico	McKinney, 1996	Jeltsch <i>et al.</i> , 1997	SOUTH AMERICA
Reynolds and Glendenning, 1949	Branscomb, 1958	McPherson <i>et al.</i> , 1988	Le Roux, 1997	Adamoli <i>et al.</i> , 1990
Savage and Swetnam, 1990	Buffington and Herbel, 1965	Nelson and Beres, 1987	Menaut <i>et al.</i> , 1990	Bücher, 1982
Smith and Schmutz, 1975	Connin <i>et al.</i> , 1997	Smeins <i>et al.</i> , 1974	Norton-Griffiths, 1979	Bücher, 1987
California	Dick-Peddle, 1993	Scanlan and Archer, 1991	O'Connor and Roux, 1995	Distel and Boo, 1996
Bossard, 1991	Gibbens <i>et al.</i> , 1992	Weltrin <i>et al.</i> , 1997	Palmer and van Rooyen, 1998	Dussart <i>et al.</i> , 1998
Bossard and Rejmanek, 1994	Hennessy <i>et al.</i> , 1983	Utah	Ramsay and Rose Innes, 1963	San José and Farías, 1983
Callaway and Davis, 1993	McCraw, 1985	Madany and West, 1983	Reid and Ellis, 1995	San José and Farías, 1991
McBride and Heady, 1968	York and Dick-Peddle, 1969	Yorks <i>et al.</i> , 1992	Ringose <i>et al.</i> , 1996	San José <i>et al.</i> , 1991
Hobbs and Mooney, 1986	Nevada	Washington	Sabiti, 1988	San José and Montes, 1997
Vivrette and Muller, 1977	Blackburn and Tueller, 1970	Rummell, 1951	Schwartz <i>et al.</i> , 1996	San José <i>et al.</i> , 1998
Williams <i>et al.</i> , 1987	North Dakota	Wyoming	Scott, 1966	Schofield and Bucher, 1986
Young and Evans, 1981	Potter and Green, 1964	Fisher <i>et al.</i> , 1987	Shantz and Turner, 1958	OTHERS
Colorado	Oregon	Regional Assessments	Skarpe, 1990	Backeus, 1992
Baker and Weisberg, 1997	Knapp and Soule, 1996	Glendenning and Paulsen, 1955	Skarpe, 1991	Binggeli, 1996
Mast <i>et al.</i> , 1997	Knapp and Soule, 1998	Gruell, 1983	Thomas and Pratt, 1967	Walker <i>et al.</i> , 1981
Mast <i>et al.</i> , 1998	Miller and Rose, 1995	Hart and Laycock, 1996	Trolope, 1982	Skarpe, 1992
Veble N and Lorenz, 1991	Miller and Halpern, 1998	Humphrey, 1958	Van Vegten, 1983	
Idaho	Miller and Rose, 1999	Humphrey, 1987	West, 1947	
Anderson and Holte, 1981	Skovlin and Thomas, 1995	Johnson, 1987	AUSTRALIA	
Burkhardt and Tisdale, 1976	Soule and Knapp, 1999	Leopold, 1951	Booth and Barker, 1981	
Zimmerman and Neunswander, 1984	Iowa	Milchunas and Lauenroth, 1993	Bowman and Panton, 1995	
	Engle <i>et al.</i> , 1996	Miller and Wigand, 1994	Bren, 1992	
	Snook, 1985	McClaran and McPherson, 1995	Brown and Carter, 1998	
			Burrows <i>et al.</i> , 1985	

Note. Documentation includes historical observations, long-term monitoring, repeat ground or aerial photography, stable carbon isotope analysis, dendrochronology, and, in some cases, simulation modeling. The focus is on arid and semi-arid "rangelands." Hence, studies documenting tree/shrub invasion of abandoned agricultural fields (cf. Smith, 1975; Johnston *et al.*, 1996; De Steven, 1991) or regeneration following forest clearing are not included here. Studies discussing or reviewing causes or consequences of woody encroachment into grasslands/savannas (cf. Humphrey, 1953; Fisher, 1977; Smeins, 1983; Rappole *et al.*, 1986; Grover and Musick, 1990; Schlesinger *et al.*, 1990; Archer *et al.*, 1994; Archer *et al.*, 1995; Idso, 1995; Polley *et al.*, 1997) are also excluded. Some papers in the list reference other papers which have documented woody plant increases in historical times (Backeus, 1992; Noble, 1997). An updated and more extensive version of this table, including a list of woody genera, can be found at <http://cnr.it.tamu.edu/rlem/faculty/archer/>.




FIGURE 2 Xerification/desertification (West, 1986) and deforestation have received much attention. Although increases in woody plant abundance in drylands are geographically widespread and well documented (Table 1), little is known of the ecological consequences of this vegetation change (adapted from Archer and Stokes, 2000).

trace gases, NO_x and aerosols from savanna fires may contribute significantly to global emissions and influence climate and atmospheric chemistry (Crutzen and Andreae, 1990; Hao et al., 1990; Crutzen and Goldammer, 1993).

Desertification has long been a topic of concern to land managers and ecologists (Moat and Hutchinson, 1995; Arnalds and Archer, 2000). More recently, changes in the storage and dynamics of C and N in the terrestrial biosphere have been evaluated with respect to deforestation, intensive agricultural practices, succession on abandoned agricultural lands, and afforestation/reforestation (Fig. 2) (Houghton et al., 1987; Post, 1993). Increased abundance of woody plants in drylands has the potential to alter land surface-atmosphere interactions and atmospheric chemistry by affecting biophysical processes, and C and N storage and dynamics (e.g., Schlesinger et al., 1990; Graetz, 1991; Bonan, 1997). Even so, its significance has yet to be thoroughly evaluated or quantified. Here, we review results from a case study of a subtropical dryland landscape which has been undergoing a transformation from grassland to savanna to woodland. Some of our recent work has explored the implications of this change in vegetation on the hydrological cycle (Brown and Archer, 1990; Midwood et al., 1998; Boutton et al., 1999). Here, our emphasis is on the rates of change in soil and plant carbon and nitrogen pools and fluxes.

3. The La Copita Case Study

3.1 Biogeographical and Historical Contexts

The La Copita Research Area (27° 40'N; 98° 12'W; elevation = 75–90 m ASL) is situated in the northeastern portion of the North American Tamaulipan Biotic Province (Blair, 1950) in the

Rio Grande Plains of southern Texas. The potential natural vegetation of this region has been classified as *Prosopis-Acacia-Andropogon-Setaria* savanna (Küchler, 1964). However, the contemporary vegetation is subtropical thorn woodland (McLendon, 1991) and occupies about 12 million ha in Texas alone (Jones, 1975). The shrubs and small trees at the study site are characteristic of dry tropical and subtropical zones in Mexico, Central America, South America (Chaco, Caatinga, Caldénal), Africa, Australia, India, and southeast Asia. In many instances, it is believed that these vegetation types have replaced grasslands over large areas since the 1800s (Table 1). Current vegetation at the La Copita site, which has been grazed by domestic livestock since the late 1800s, consists of savanna parklands in sandy loam uplands that grade into closed-canopy woodlands in clay loam lowland drainages. All wooded landscape elements (upland shrub clusters and groves; lowland playa and drainage woodlands) are typically dominated by the leguminous tree *Prosopis glandulosa* in the overstory, with an understory mixture of evergreen, winter-deciduous, and summer-deciduous shrubs. Climate of the region is subtropical (mean annual temperature 22.4 °C) with warm, moist winters and hot, dry summers. Mean annual rainfall is 720 mm and highly variable (CV = 35%).

Reports from settlers indicate that much of southern Texas was grassland or open savanna in the mid-1800s (Inglis, 1964). Historical aerial photography demonstrates that woody plant cover on La Copita increased from 10% in 1941 to 40% in 1983 (Archer et al., 1988). $\delta^{13}\text{C}$ and radiocarbon analyses of soil organic carbon have confirmed that C_3 trees and shrubs have displaced C_4 grasses in upland and lowland portions of the landscape in the past 100 years (Boutton et al., 1998). Plant growth (Archer, 1989) and transition probability models (Scanlan and Archer, 1991), substantiated by tree ring analysis (Boutton et al., 1998), indicate that most trees on the site have established over the past 100 years. The successional processes involved in woody plant community development and topographic controls over spatial patterns of tree/shrub expansion have been elucidated (Archer, 1995b). Armed with information from these prior studies, we are now poised to ascertain the biogeochemical consequences of succession from grassland to woodland.

3.2 Herbaceous Retrogression and Soil Carbon Losses

Simulations with the CENTURY biogeochemistry model (Parton et al., 1987, 1988, 1993) parameterized for assumed presettlement conditions ("light" grazing, fire at 10-year intervals), soil texture, and climate of the La Copita projected that soil organic carbon (SOC) would have been on the order of 2500 g m⁻² to a depth of 0–20 cm (Hibbard, 1995). The SOC values from this assessment were then used as a baseline against which historic effects of heavy, continuous livestock grazing were evaluated. In a subsequent model run, intensification of grazing and removal of fire were initiated in 1850, a date approximating the advent of widespread, unregulated livestock grazing in southern Texas (Lehman,




FIGURE 3 Simulation model reconstruction of changes in soil organic carbon (0–20 cm) on an upland, sandy loam soil and a lowland, clay loam soil at the La Copita Research Area in southern Texas, USA (from Hibbard, 1995). Solid lines depict steady-state SOC values expected for the climate of the site under light grazing and fire every 10 years (presettlement conditions). Dashed lines depict changes in SOC predicted to occur on two major soil types after the onset of heavy, continuous livestock grazing and cessation of fire. Steady-state values for heavy, continuous grazing and cessation of fire are within 5% of field measurements on present-day grasslands.

1969). This simulation produced a 16–29% reduction in SOC of sandy loam upland and clay loam lowland soils, respectively (Fig. 3). These results appear reasonable in that the model-generated steady-state SOC levels for sandy loam uplands (2062 g m^{-2}) approximated the average pool sizes measured in present-day grassland communities on these soils (2087 g m^{-2}). Further, the grazing-induced reductions in SOC predicted by the simulation were comparable to field measurements reported for other grazed grasslands (Bauer *et al.*, 1987; Frank *et al.*, 1995; but see also Milchunas and Lauenroth 1993). CENTURY simulations did not explicitly include potential erosion losses. The fact that simulated historic changes in SOC approximated those currently observed at the site therefore suggests that such losses may have been minimal. This inference seems reasonable, since the La Copita landscapes have relatively little topographic relief (1–3% slopes) and show no obvious physical signs of erosion (pedestals, rills, gulleys). Furthermore, soil profile structure in low-lying portions of the landscape shows no pedogenic evidence of significant translocation of soils from uplands. Elevated C and N pools in soils of developing woody communities (summarized later) thus appear to be the result of *in situ* accumulations induced by trees and shrubs rather than losses from grazed grasslands.

The present-day herbaceous vegetation is dominated by a low cover of ephemeral dicots and short-statured, weakly perennial grasses. In contrast, herbaceous vegetation on relict, protected grasslands in the region is characterized by mid- to tall-statured perennial grasses whose potential productivity ($500\text{--}600 \text{ g m}^{-2}$; SCS, 1979) is two to three times that which has been recorded at La Copita ($<270 \text{ g m}^{-2}$; Vega, 1991; Hibbard, 1995). Thus, it is reasonable to conclude that soil C and N storage has declined in

herbaceous communities over the past century as a result of changes in species composition, microclimate, and biomass production attributable to heavy, continuous livestock grazing on this site.

3.3 Woody Plant Encroachment and Ecosystem Biogeochemistry

Changes in soils and microclimate accompanying long-term heavy grazing may have shifted the balance in favor of N_2 -fixing or evergreen woody plants which are better adapted than grasses to nutrient-poor soils and warmer, drier microenvironments. The establishment of trees and shrubs would have been further augmented by grazing-induced reductions in herbaceous competition and fire (Archer, 1995a). In addition, the woody plants at La Copita are highly unpalatable, and browsing by wildlife or cattle is minimal. However, fruits of the dominant tree invader (*Prosopis glandulosa*) are readily consumed by livestock, which disperse large numbers of viable seeds into grasslands (Brown and Archer, 1987). Thus, heavy, continuous, and preferential grazing of grasses by livestock has promoted woody plant encroachment via numerous direct and indirect effects (Archer, 1994). As woody communities develop in grazed grasslands, plant and soil C and N pool sizes and flux rates change as described in the following sections.

3.3.1 Plant Carbon Pools

Quantitative changes in woody plant cover at La Copita are depicted in Fig. 4 and 5. To ascertain the effects of these vegetation changes on plant carbon stocks, we linked CENTURY with a plant succession model developed for La Copita (Scanlan and Archer, 1991). We initiated woody plant encroachment in the late 1800s on a heavily grazed, fire-free landscape in which SOC content had been reduced by grazing (Fig. 3). The landscape, consisting of a




FIGURE 4 Changes in total woody plant cover on three replicated landscapes at the La Copita site in southern Texas. See Figure 5 for spatial pattern of change in various patch types on Landscape 1 (Archer and Boutton, unpublished).




FIGURE 5 Landscape-scale changes in herbaceous and woody plant community cover from 1950 to 1990 in upland (herbaceous, discrete cluster, and grove) and lowland (woodland) plant communities at the La Copita site in southern Texas (Archer and Boutton, unpublished). Values given below dates are total hectares of woody cover (cluster + grove + woodland) for the 11.06 ha "pixel." See Figure 4 (Landscape 1) for changes in percentage woody cover.

sandy loam upland and a clay loam intermittent drainage, was populated with woody plants based on rainfall-driven transition probabilities. Grassland and forest CENTURY subroutines were applied, as appropriate, to compute plant carbon and SOC (0–20 cm) stocks in wooded and herbaceous landscape elements. At each time step, plant carbon and SOC were estimated by summing across the entire landscape (upland plus lowland vegetation patch types). Results suggest that the development of the present-day savanna parkland–woodland complex has increased plant carbon stocks 10-fold over that which would be present had the "pristine" grassland vegetation been maintained on the site (Fig. 6). Part of that increase is attributable to an increase in aboveground net pri-

mary productivity (Table 2) and part of it represents the decline in tissue turnover which occurs when herbaceous vegetation is replaced by woody vegetation. These results are conservative in that CENTURY simulations include root C mass only in the top 20 cm of soil. Biomass distributions of woody plant roots at La Copita (Watts, 1993; Boutton *et al.*, 1998; 1999; Midwood *et al.*, 1998; Gill and Burke, 1999) are typical of those of other dryland tree/shrub systems (Jackson *et al.*, 1996; Canadell *et al.*, 1996) where, relative to grasslands, there is substantially greater mass at deeper depths where turnover and decomposition are likely to be reduced. The fact that fluctuations in monthly woody plant root biomass in upper soil horizons exceeded monthly foliar litter inputs by one to

TABLE 2 Contrasts in Aboveground Net Primary Production (ANPP), Soil Physical Properties, Organic Carbon and Total Nitrogen Pools (0- to 10-cm depth), and Fluxes in Soils Associated with Woody Plant and Grazed Grassland Communities in a Sandy Loam Upland Landscape at the La Copita Research Area in Southern Texas

Parameter	Community Type	
	Herbaceous	Woody Plant
ANPP ($\text{Mg ha}^{-1} \text{ year}^{-1}$) ^a	1.9 – 3.4	5.1 – 6.0
Bulk density (g cm^{-3}) ^a	1.4 ± 0.01	1.1 ± 0.04
% Clay ^a	20 ± 0.7	20 ± 1
Fine roots (g m^{-2}) ^a	100 – 175	400 – 700
Coarse roots (g m^{-2}) ^a	100 – 400	400 – 1,100
Organic C, % g m^{-2}	0.84 ± 0.05 1165 ± 67	2.2 ± 0.23 2352 ± 276
Potential C mineralization ($\text{mg C kg}^{-1} \text{ soil day}^{-1}$) ^b	7.3 ± 5.7	15.5 ± 6.8
Soil respiration ($\text{mg CO}_2 \text{ m}^{-2} \text{ year}^{-1}$) ^b	611 ± 83	730 ± 67
Q_{10} values for <i>in situ</i> soil respiration ^b	1.2	1.4 ± 2.7
Total N, % g m^{-2}	0.07 ± 0.00 91 ± 6	0.18 ± 0.02 192 ± 20
N mineralization ^a , $\text{g N m}^{-2} \text{ year}^{-1}$	6 ± 1	22 ± 2
$\mu\text{g N g}^{-1} \text{ year}^{-1}$	42 ± 5	200 ± 18
NO flux ($\text{ng NO-N cm}^{-2} \text{ h}^{-1}$) ^c		
Dry soil	0.2 ± 0.07	2.8 ± 0.25
Wet soil	1.1 ± 0.11	16.2 ± 2.03

Note. Maximum and minimum monthly values for samples obtained over an annual cycle are shown for root-standing crop (coarse roots ≥ 0.1 -mm diameter); a range is presented for ANPP and Q_{10} . All other values are means \pm SE.

^aHibbard, 1995.

^bMcCulley, 1998.

^cCole et al. 1996.

two orders of magnitude (Table 2) suggests that belowground inputs of organic matter drive changes in soil physical and chemical properties subsequent to woody plant establishment in grasslands. These substantial fluctuations in woody plant root biomass suggest a high turnover, which is consistent with detailed observations on woody plants in other systems (Eissenstat and Yanai, 1997; Hendricks *et al.*, 1997). In addition, turnover of grass roots may be slower than has been generally assumed (Milchunas *et al.*, 1992). Thus, increases in aboveground and belowground net primary productivities may accompany woody plant encroachment into grasslands and foster C and N accumulation.

As an independent test of the reconstruction in Figure 6, we quantified aboveground plant carbon density in patches representing the dominant community types at La Copita. This was accomplished using allometric relationships and belt-transect surveys (Northup *et al.*, 1996). Plant carbon density was then multiplied by community area measured on aerial photographs (1950, 1976, 1990) to obtain a community-level estimate. Estimates for each community type were then summed to obtain landscape-scale estimates. For patches representing various woody and herbaceous community types, CENTURY estimates of aboveground carbon density were lower than field-based estimates (Table 3), further suggesting that model estimates were conservative. Aboveground carbon density differed substantially among




FIGURE 6 Modeled changes in whole-landscape (upland + lowland and all patch types therein) plant carbon density (aboveground + roots to 20 cm) accompanying succession from grassland to savanna parkland/woodland (from Hibbard, 1995). Dashed line depicts steady-state expected for a lightly grazed grassland landscape (upland + lowland communities pooled) with fire at 10-year intervals and no woody plants; solid line depicts steady-state plant carbon density for heavily, continuously grazed grassland landscape with no fire and no woody plants. Changes in woody plant abundance on each soil type were directed by a succession model (Scanlan and Archer, 1991); subsequent changes in plant carbon stocks were then assessed with a biogeochemistry model (CENTURY; Parton *et al.*, 1994). See Figure 7 for validation results.

the three landscapes inventoried in 1950 (Fig. 7), primarily reflecting differences in woody plant cover on this date (Fig. 4). By 1990, woody cover and carbon density were comparable on the three landscapes. The CENTURY-succession model estimates of aboveground carbon density for an "average" landscape in 1950, 1976, and 1990 closely approximated those obtained from the field-historical aerial photo approach.

3.3.2. Nonmethane Hydrocarbon Fluxes

On a regional basis, shifts from grass to woody plant domination have the potential to influence biophysical aspects of land-atmosphere interactions, such as albedo, evapotranspiration, boundary layer conditions, and dust loading (e.g., Bryant *et al.*,

TABLE 3 Observed and Predicted Aboveground Carbon Density in Patches Representing Tree/Shrub and Grassland Communities at La Copita, Texas

Topoedaphic Setting	Patch Type	Carbon Density (kg m^{-2})	
		Field Estimate	Model Estimate
Sandy loam upland	Cluster	2.9 ± 0.4	2.2
	Grove	6.3 ± 0.8	4.0
	Grassland ^a	0.05 ± 0.00	0.04
Clay loam lowland	Woodland	5.8 ± 0.8	4.5

Note. Observed data (means \pm SE) are based on belt transects and plant size-biomass relationships for woody communities (Archer and Boutton, unpublished) and on clipped plots in grasslands (Archer, unpublished). Predicted values are CENTURY estimates for 100-year-old patches (Hibbard, 1995).

^aPeak aboveground biomass.




FIGURE 7 Changes in aboveground plant carbon density on three landscapes at the La Copita from 1950 to 1990 (Archer and Boutton, unpublished). Patch/soil-specific field estimates of plant carbon density (Northup *et al.*, 1996, McMurtry, Nelson and Archer, unpublished) were multiplied by patch area as measured in aerial photographs to generate whole-landscape estimates. Dashed lines denote predictions from linked CENTURY-succession model (Hibbard, 1995). See Figure 4 for changes in woody cover on the three landscapes.

1990; Pilke and Avissar, 1990; Graetz, 1991). These changes in vegetation may also influence atmospheric oxidizing capacity, aerosol burden, and radiative properties by affecting emissions of non-methane hydrocarbons (NMHCs) such as terpenes, isoprene, and other aromatics (Fehsenfeld *et al.*, 1992). There are many sources of atmospheric NMHCs, but >90% of the global annual emission is from vegetation (Guenther *et al.*, 1995). NMHC emissions are therefore highly dependent on species composition as constrained by environmental conditions which influence plant physiology and production. The high temperatures and solar radiation fluxes associated with subtropical and tropical grasslands and savannas make these geographically extensive bioclimatic regions large potential sources of biogenic NMHC emissions. However, grasses are typically low emitters of NMHCs, whereas emissions from trees and shrubs in forest systems are highly variable, with some species being low emitters and other species being high emitters.

We hypothesized that foliar emissions of NMHCs in woody plants would be positively correlated with leaf longevity and inversely related to photosynthetic capacity. Plants characterized by low photosynthetic capacities and slow growth rates (e.g., evergreens) depend on extended leaf longevity to achieve a positive carbon balance. Preferential allocation to secondary compounds such as terpenes would help ensure foliage longevity by reducing levels of herbivory. Species with low photosynthetic capacities and high levels of secondary compounds should also dominate understory environments where low light levels preclude high growth rates and where plants are more accessible to browsers. In contrast, species selected for competitive ability would have high photosynthetic rates, high growth rates, and high rates of tissue turnover (e.g., deciduous shrubs). Allocation to secondary compounds that deter herbivory would be of lower priority since leaf

longevity is less critical to realizing a positive return on foliar investments. Such plants would be expected to preferentially allocate resources such as nitrogen to the carboxylating enzyme and productive tissues rather than to structural tissues or secondary compounds such as terpenes, and would thus be low NMHC emitters (or isoprene emitters, since isoprene is not known to be associated with defense; Coley *et al.*, 1985).

To test these hypotheses, we screened plant species representing the major growth forms at La Copita for NMHC emissions. As expected, grasses had low NMHC emission rates and several common woody species had high emission rates (Guenther *et al.*, 1999). However, there was little evidence of emissions being consistently related to woody plant taxonomy, growth form, or functional group. As a result, generalizations regarding NMHC emissions spectra for tree/shrub species assemblages in other systems do not appear feasible.

To determine if biogenic NMHC emissions have been altered as a result of the change in land cover from grass to woody plant domination at La Copita, a vegetation change model (Scanlan and Archer, 1991) was then linked with a model which predicted NMHC emissions as a function of foliar density, leaf temperature, and photosynthetic photon flux density as modulated by ambient temperature, cloud cover, precipitation, relative humidity, and wind speed (Guenther *et al.*, 1995; Guenther, 1997).

Linkage of the biogenic emissions model with the plant succession model indicated that land cover change since the early 1800s has elicited a threefold increase in isoprene emissions (Fig. 8). This increase reflected changes in vegetation composition and increases in foliar density. Model predictions of current NMHC emissions were within 20% of those measured by a tower flux system. Detailed field measurements on two common shrub species indicated that isoprene emission increased exponentially with increases in leaf temperature from 20 to 40°C and were not suppressed by drought stress. Accordingly, the model predicted that under a projected 2X-CO₂ climate, present-day biogenic NMHC emissions would double.

These estimates of changes in NMHC emissions associated with the conversion of grassland to woodland are in accordance with estimates in other ecosystems. For example, Klinger *et al.* (1998) documented a fourfold increase in total terpenoid emissions per unit foliar mass along a savanna to woodland transect in Central Africa. These changes in NMHC emissions associated with vegetation change in subtropical Texas and tropical Africa also mirror those reported for temperate forest (Martin and Guenther, 1995). Together, these results indicate the magnitude of change in NMHC emissions that could occur when climate and vegetation composition are altered. The importance of these increases in NMHC emissions is magnified at La Copita, as they occur in conjunction with elevated nitric oxide (NO) emissions from shrub-modified soils (Table 2; see Sec. 3.3.4 for elaboration).

Why are vegetation-induced increases in NMHC of concern? Biogenic hydrocarbons play an important role in generating pollutants such as O₃, CO, and organic peroxides, while influencing hydroxyl radical (OH⁻) chemistry to reduce atmospheric oxida-




FIGURE 8 Changes in nonmethane hydrocarbon (isoprene) emissions predicted to accompany a shift from savanna grassland to a savanna woodland at the La Copita site in southern Texas (based on Guenther *et al.*, 1999). Predictions from a coupled succession–NMHC emission model are compared with values measured from flux towers. The “measured” values shown for the historic landscape are from a tower located in a savanna grassland landscape with low woody cover.

tion capacity and increase the residence time of greenhouse gases. It has been estimated that to meet current air quality standards for tropospheric ozone, anthropogenic hydrocarbon emissions would have to be reduced by only 30% in the absence of natural isoprene emissions, but by 70% in the presence of them (Monson *et al.*, 1991). Changes in NMHC–NO emissions associated with regional conversion of grassland to shrubland may therefore constitute a “moving baseline” from which to gauge tropospheric ozone production triggered by emissions from automobiles or industrial sources.

3.3.3 Soil C and N Pools

Once established, woody plants alter soils and microclimate in their immediate vicinity to affect both pool sizes and flux rates of nutrients. The result is the formation of “islands of fertility,” a phenomenon which has been widely quantified in drylands (see Charley and West, 1975; Schlesinger *et al.*, 1990; Scholes and Archer, 1997; special issue of *Biogeochemistry* 42 (1/2) 1998). Three general mechanisms have been proposed to account for this (e.g., Virginia, 1986): (1) woody plants act as nutrient pumps, drawing nutrients from deep soil horizons and laterally from areas beyond the canopy, depositing them beneath the canopy via stem flow, litterfall, and canopy leaching; (2) tall, aerodynamically rough woody plant canopies trap nutrient-laden atmospheric dust that rain washes off the leaves and into the subcanopy soil; and (3) woody plants may serve as focal points attracting roosting birds, insects, and mammals seeking food, shade, or cover. These animals may enrich the soil via defecation and burrowing. For these reasons, soil carbon and nitrogen pools should increase subsequent to woody plant colonization in grazed grasslands.

At La Copita, surficial (0–10 cm) soils associated with woody plants known to have encroached over the past century have a lower bulk density, contain more root biomass, have higher concentrations of SOC and total N, and have greater rates of respiration and N mineralization than soils associated with the remaining grazed grassland communities (Table 2). As the continuity of woody plant cover increases through time, the landscape-scale soil nutrient pools and fluxes would be expected to increase and become more homogeneously distributed. Accordingly, the linked CENTURY–succession model exercise (see Section 3.3.1) predicted that by 1950, landscape-scale SOC had returned to levels which would have occurred had the “pristine” grasslands been maintained on the site (Fig. 9). By the early 1990s, landscape-scale SOC levels were about 10% higher than those expected for the “pristine” grassland, and about 30% higher than those for a heavily grazed grassland not experiencing woody plant encroachment. Forward model projections suggest SOC aggradation will continue for several hundred years, reaching equilibrium levels three times those of the present-day grazed grassland communities.

While the “island of fertility” phenomenon has been widely recognized, little is known of the rates of nutrient enrichment in tree-dominated patches. Total C and N in soil under *Acacia senegal* and *Balanites aegyptiaca* tree canopies were positively correlated with tree girth ($r^2 = 0.62$ and 0.71 , respectively; Bernhard-Reversat, 1982), indicating net accumulation with time of woody plant occupancy of a patch. In temperate old fields undergoing forest succession, carbon storage increased 40% in plant + soil pools over 40 years (Johnston *et al.*, 1996). At La Copita, soil C and N were quantified under *Prosopis glandulosa* trees whose age was determined by annual ring counts. Soil organic carbon storage (top 20 cm of soil) increased linearly with tree stem age at




FIGURE 9 Changes in soil organic carbon (SOC; 0–20 cm) predicted to accompany woody plant encroachment into a grazed landscape consisting of a sandy loam uplands and clay loam intermittent drainages at the La Copita site (from Hibbard, 1995). Dashed line depicts steady-state SOC expected for a lightly grazed grassland landscape (upland + lowland communities pooled) with fire at 10-year intervals and no woody plants; solid line depicts steady-state SOC for heavily, continuously grazed grassland landscape with no fire and no woody plants (see Fig. 3). Bars denote SOC summed across the entire landscape and include both grassland and woody plant communities. Note that by 1950, SOC levels had increased to a level comparable to that of the “pristine” grassland (cross-hatched bar). Changes in woody plant abundance on each soil type were directed by a succession model (Scanlan and Archer, 1991); subsequent changes in soil carbon were then assessed with a biogeochemistry model (CENTURY; Parton *et al.*, 1994).

rates ranging from 11.8 to 21.5 g C m⁻² year⁻¹ in sandy loam uplands woody patch types to 47.2 g C m⁻² year⁻¹ in moister, clay loam woodland patches (Table 4). Rates of total N accumulation (top 20 cm of soil) ranged from 1.9 to 2.7 g N m⁻² year⁻¹ in sandy uplands and averaged 4.6 g N m⁻² year⁻¹ in clay loam lowlands. However, woody plant age explained only 21–68% of the variation in soil C and N sequestration rates. These low *r*² values may indicate that tree stem ages do not accurately reflect plant ages, possibly due to past disturbance and subsequent vegetative regeneration of woody cover. Low *r*²s may also indicate that factors un-

related to time of tree occupancy influence soil C and N under tree canopies. Such factors may include small-scale heterogeneity associated with large mammal or bird defecation, soil mixing by small mammals and arthropods, or patch-specific differences in the species composition, productivity, and rate of development of the understory community.

Modeling experiments allowed us to control for factors that might cause variation in field-based estimates of woody plant age–SOC relationships. Model estimates of SOC accumulation were comparable to field estimates for upland patch types and substantially lower than field estimates for lowland patch types (Table 4). Model estimates of soil N accumulation were substantially lower than field estimates, especially in lowlands. Given that woody patch age explained only 26–68% of the variance in soil C and N content, our field estimates of accumulation rates cannot be taken as definitive. Model results underestimated field observations, especially for N. Reliability of model estimates of soil carbon could likely be improved with a better understanding of how turnover of the substantial root mass (Table 2) might differ among patch types. Model estimates of soil N are likely constrained by lack of information on inputs associated with N₂ fixation, atmospheric N deposition, translocation between uplands and lowlands, and root turnover.

3.3.4 Soil C and N Dynamics

Increases in the C and N pools of soils associated with woody plant communities developing on grazed grasslands at La Copita have been accompanied by increases in soil respiration, N mineralization, and nitric oxide (NO) emissions (Table 2). The increase in NO fluxes accompanying expansion of woody plants into grasslands at La Copita is noteworthy. Nitric oxide plays several critical roles in atmospheric chemistry by contributing to acid rain and by catalyzing the formation of photochemical smog and tropospheric ozone. The latter is potentially accentuated in the La Copita setting, since NO and hydrocarbon emissions (see Sec. 3.3.2) are concomitantly elevated subsequent to woody plant establishment.

The quality and quantity of organic matter inputs interact to drive soil metabolic activity (Zak *et al.*, 1994). Hence, annual soil

TABLE 4 Estimated Rates of Organic Carbon and Total Nitrogen Accumulation in Soils (0- to 20-cm Depth) Developing Beneath Woody Plants Establishing on a Former Grassland

Location	Soil Texture	Patch Type	g C m ⁻² year ⁻¹		g N m ⁻² year ⁻¹	
			Field	Modeled	Field	Modeled
Upland	Sandy loam	Shrub Cluster	21.5 (<i>r</i> ² = 0.26)	18.5	2.67 (<i>r</i> ² = 0.45)	1.15
		Grove	11.8 (<i>r</i> ² = 0.21)	10.5	1.90 (<i>r</i> ² = 0.51)	0.87
		Woodland	47.2 (<i>r</i> ² = 0.57)	13.1	4.64 (<i>r</i> ² = 0.68)	0.58

Note. Field data are from linear correlations between patch age (determined by dendrochronology) and soil C and N mass (Boutton and Archer, unpublished). Model estimates are from CENTURY simulations (Hibbard, 1995). Descriptions of contrasting woody patch types can be found in Archer (1995b).

respiration rates are positively correlated with net primary productivity (Raich and Schlesinger, 1992). The elevated carbon fluxes observed with the development of woody communities in semiarid La Copita grasslands may reflect increased root (Table 2) and leaf biomass inputs and enhancement of soil moisture beneath woody plant canopies (via concentration of rainfall from stem flow, hydraulic lift, and/or reduced evaporation). Together, these biotic and abiotic factors may interact to stimulate microbial activity relative to that in grass-dominated soils. In fact, microbial biomass in woody communities is comparable to or higher than that in grassland communities at La Copita (McCulley, 1998). However, experimental irrigation, which alleviated plant water stress, enhanced photosynthesis (McMurtry, 1997), and increased soil respiration, elicited a decrease in soil microbial biomass. This suggests the elevated soil respiration observed in woody plant communities at La Copita may be a consequence more of changes in root biomass (Table 2) and respiration than of changes in microbial biomass and activity.

To estimate landscape-scale changes in soil CO₂ flux, we multiplied patch/soil-specific estimates of annual soil respiration (McCulley, 1998) by patch area. We then computed changes in patch area with a succession model (Scanlan and Archer, 1991). Landscape-scale soil respiration (kg C ha⁻¹ year⁻¹) is projected to have increased from 6687 (200 YBP) to 7377 (1990s) to 7602 (200 years in future)(Table 5). This represents a 10.3% increase with the transition from historic grassland savanna to the present-day savanna parkland-thorn woodland complex, with an additional 3% increase occurring if the present savanna parkland progresses

to woodland. If mean annual temperatures increase as projected in general circulation models, further increases in soil respiration would be expected (all other factors being equal). Indeed, Q₁₀ values for soil respiration in woody plant communities (1.4, 2.7, and 2.3 in cluster, grove, and woodland types, respectively) exceed those of grazed grasslands (1.2) at La Copita (McCulley, 1998). This suggests that if future temperature changes occur, the importance of recent and projected future vegetation changes on soil respiration will be further magnified. For example, the magnitude of increase in soil respiration from past grassland savanna with MAT of 22.4°C to future woodland with MAT of 28.4°C would be 22.5% (= 6687–8197 kg C ha⁻¹ year⁻¹; based on Raich and Schlesinger, 1992) to 99.3% (= 6687–13,328 kg C ha⁻¹ year⁻¹; based on McCulley, 1998)(Table 5). Potential changes in the amount, seasonality, and effectiveness of rainfall would have important, but as yet unknown, effects on these projections.

3.3.5 Soils as Sources and Sinks

Elevated fluxes of C and N from plants and soils following grassland-to-woodland conversion at La Copita suggest a potential for augmenting greenhouse gas accumulation and altering tropospheric chemistry, particularly if woody plant encroachment has been geographically widespread (as suggested in Table 1). However, as noted in Sec. 3.3.3, organic C and N have accumulated in soils of developing woody plant communities at La Copita, despite elevated fluxes and higher turnover rates. This indicates that inputs have exceeded outputs and that soils and vegetation at La

TABLE 5 Projected Landscape-Scale Changes in Annual Soil Respiration (SR; kg C ha⁻¹ year⁻¹) Accompanying Succession from an Open Savanna/Grassland to Woodland and Potential Changes in Mean Annual Temperature

Landscape-Scale Soil Respiration (kg C ha ⁻¹ year ⁻¹)			
Mean Annual Temperature (°C)	Past Grassland (200 YBP) ^a	Present Savanna Parkland/Woodland Complex	Future Thorn Woodland (200 YAP) ^b
A. Based on MAT/MAP regression in Raich and Schlesinger (1992)			
22.4	6687	7377	7602
25.4	6948	7666	7899
28.4	7209	7954	8197
B. Based on Q ₁₀ values from McCulley (1998)			
25.4	8083	9938	10,465
28.4	9480	12,499	13,328

Note. Patch- (grass and various woody communities) and soil-specific SR rates measured monthly over an annual cycle at La Copita (McCulley, 1998) were multiplied by the area of respective community types (Scanlan and Archer, 1991). Effects of mean annual temperature change (MAT, °C) on SR were estimated from (A) equations in Raich and Schlesinger (1992); for La Copita (MAT = 22.4°C and MAP = 720 mm) a 3 and 6°C increase in MAT would produce a 3.9 and 7.8% increase, respectively, in soil respiration; and (B) Q₁₀ values of *in situ*, community-specific soil respiration from McCulley (1998). Estimates are probably conservative, as respiration rates used in computations were measured during a below-normal rainfall year.

^aYBP, years before present.

^bYAP, years after present.

Copita have been functioning as C and N sinks over the past century. A variety of factors might interact to account for the observed increases in soil C and N pools:

- The trees and shrubs which have displaced grasses may be more productive aboveground and belowground and hence deliver more organic matter into soils (see root biomass and ANPP in Table 2).
- Leaves of leguminous and nonleguminous woody plants at La Copita have higher [N] than grasses (2–4% vs <1%; Archer, unpublished). However, woody plants in these landscapes are seldom browsed by livestock or wildlife, suggesting high concentrations of secondary compounds. This could result in a significant litter quality \times quantity interaction, whereby
 - a large fraction of the foliar biomass produced by trees and shrubs goes into the soil pool and directly as litter rather than through the herbivory pathway, and
 - a larger fraction of foliar biomass inputs from woody plants may be resistant to decomposition.
- Woody litter inputs and the coarser, more lignified roots of shrubs would promote C and N accumulation compared to that of grass roots and shoots.
- Shading by tree/shrub canopies reduces soil temperatures relative to those in grassland (Archer, 1995b), thus constraining potential mineralization (Q_{10} effect).
- Nitrogen accumulation is potentially a consequence of N_2 fixation by leguminous shrubs common to the site (*P. glandulosa* and several *Acacia* spp.) and/or the uptake and lateral translocation of N from grassland patches. While nodulation has been induced in controlled environments and observed under field conditions at the La Copita site (Zitzer et al., 1996) and elsewhere (Virginia et al., 1986; Johnson and Mayeux, 1990), methodological constraints have prevented quantification of N_2 fixation (Handley and Scrimgeour, 1997; Liao et al., 1999). Root distribution studies (Watts, 1993) discount the lateral foraging hypothesis.
- La Copita is within ca. 70 km of a major oil refinery center (Corpus Christi, TX) and atmospheric N deposition has likely been significant over the past 50–75 years (e.g., Holland et al., 1999). Increased N availability may have promoted woody plant expansion (e.g., Köchy, 1999) by alleviating grass–woody plant competition for soil N and by promoting growth of woody plants more than that of grazed grasses. This, in turn, may have translated into greater organic C and N inputs into soils associated with woody plants.

3.3.6 An Uncertain Future

Prosopis glandulosa currently dominates the overstory in upland and lowland woody plant communities. Depending on patch type, it constitutes 40–90% of the above-ground biomass (Archer and Boutton, unpublished) and 30–70% of the coarse root (>1-mm diameter) biomass (Watts, 1993). As such, the dynamics of *P. glandulosa*

must be a primary driver of changes in plant and soil C and N stocks at La Copita. Future increases in landscape nutrient pools and fluxes will reflect a combination of (a) continued growth of *P. glandulosa* and associated shrubs in existing woody plant communities and (b) expansion of woody plants into the remaining grasslands.

How likely is continued expansion? That may depend on land management practices. Relaxation of grazing pressure could enable grass biomass to accumulate and fire (prescribed or natural) to occur. Together, these could retard expansion and growth of woody plants. However, the La Copita appears to have crossed a threshold, whereby soils, seed banks, and vegetative regenerative characteristics are such that reductions in grazing pressure may be of little consequence (Archer, 1996). Relaxation of grazing would influence woody plant establishment in grassland primarily through its influences on the fire regime (Archer, 1995a; Brown and Archer, 1999). However, the remaining herbaceous clearings are small and discontinuously distributed. Hence, even if fine fuels were to accumulate, fires would be highly localized. Such fires might prevent future encroachment into remaining grassland clearings but would not likely convert woody plant communities to grassland, since the trees and shrubs at La Copita quickly regenerate by sprouting after disturbance (Scanlan, 1988; Flinn et al., 1992). Expenses for clearing woody vegetation via mechanical or chemical treatments are prohibitive and generally not cost-effective, especially since the effects of the treatments are relatively short-lived. Thus, the likelihood of continued woody plant dominance is high, even with aggressive land management practices which might favor grasses.

The succession model which simulates the expansion of woody plants into remaining grasslands (Scanlan and Archer, 1991) projects that with heavy grazing and no fire, woody cover will continue to increase until the landscape goes to nearly complete canopy closure. This assumption has been substantiated by field data which indicate extension of lateral roots beyond woody canopies is minimal (Watts, 1993). Hence, there is little opportunity for between-cluster root competition and density-dependent regulation. As a result, tree/shrub densities may continue to increase until all available herbaceous clearings have been occupied and canopy cover is nearly continuous. Accordingly, woody patches on contrasting upland soils and woody patches on uplands that border woody communities of lowlands have grown and coalesced from the 1940s through the 1990s (Archer et al., 1988; Stokes, 1999). However, recent studies suggest that La Copita landscapes may be reaching their carrying capacity for woody plants, owing to topodaphic constraints (Stroh, 1995; Stokes, 1999). If this is the case, future changes in C and N pools will occur only with growth of plants in existing woody communities. Only time will tell if this is indeed the case.

As the current population of the dominant *P. glandulosa* ages, growth and biomass accumulation rates should slow, unless other woody species compensate. The understory shrubs that colonize beneath the *Prosopis* canopy subsequent to its establishment in grasslands slow *Prosopis* growth and seed production, hasten its

mortality (Barnes and Archer, 1998) and prevent its reestablishment (Archer, 1995b). Thus, it appears that *P. glandulosa* will not be a component of future woodlands on this landscape. Assessments to date suggest that over the short-term, loss of *Prosopis* will not adversely affect understory shrub productivity or soil C and N pools (Hibbard, 1995; Barnes and Archer, 1996). However, none of the associated woody species appears to have the genetic potential to achieve the size of mature *Prosopis* plants, either above- or below-ground. Thus, there may be less potential for carbon storage once *Prosopis* is lost from the system, unless the remaining understory species compensate by increasing their productivity. In addition, the carbon currently stored in *Prosopis* biomass would be lost via death and decomposition, albeit rather slowly. It would be interesting to explore these scenarios with a linked CENTURY-succession model. Unfortunately, we know little of the productivity of the understory shrubs. Further, the maximum age of *P. glandulosa* is unknown and we have little basis on which to prescribe mortality from the present-day population.

4. Degradation: Ecological Versus Socioeconomic

Degradation associated with "desertification" or "xerification" in arid environments (West, 1986; Rapport and Whitford, 1999) or "deforestation" in humid environments is in sharp contrast to that associated with "thicketization" of grasslands and savannas in mesic environments (Fig. 2). Desertification and deforestation typically have negative consequences both ecologically and socioeconomically. "Thicketization" has some adverse socioeconomic implications, as it reduces the capacity of rangelands for subsistence or commercial livestock production. However, it does not necessarily represent a degraded system with respect to biodiversity, productivity, nutrient cycling, and other important ecological characteristics.

Today's La Copita landscape is clearly different from that of 100–200 YBP, but is it "degraded"? The conceptual model in Figure 10, based on the La Copita case study, proposes a degradation phase (Fig. 3) followed by an aggradation phase (Figs. 6 and 9, Table 2) that begins when unpalatable woody plants establish, grow, modify microclimate, and enrich soil nutrients. Present-day landscapes at La Copita are a rich mosaic of productive woodlands and tree-shrub patches interspersed with remnant grass-dominated patches. Current plant and soil C and N mass is substantially greater than that which occurred under "pristine" conditions. In addition, these landscapes are highly resilient following disturbance (Scanlan, 1988; Flinn *et al.*, 1992) and provide habitat for numerous wildlife species, both game and nongame. So, in this case, the system that has developed following an initial degradation phase is now ecologically diverse, productive, and functional. It would seem that it is "degraded" or "dysfunctional" (Tongway and Ludwig, 1997) only with respect to its socioeconomic value for cattle grazing. However, it has other potential socioeconomic values whose realization would necessitate a change from traditional land




FIGURE 10 Conceptual model of ecosystem changes accompanying grazing-induced succession from grassland to woodland based on the La Copita case study. Dashed lines depict hypothesized upper and lower bounds of Y values; (i.e., lower bound is that which might occur in absence of woody plant encroachment (Fig. 3)); upper bound is that which might occur when woody plant communities mature (e.g., forward projection of Figs. 6 and 9). Values for biodiversity will vary substantially from system to system; in some cases (e.g., *Juniperus* systems) low-diversity monocultures of woody vegetation may develop. At the La Copita, this degradation-aggradation cycle has occurred over ca. 100–150 years.

uses. These include alternative classes of livestock (e.g., goats), lease hunting, charcoal production, and ecotourism.

Given the demonstrated potential for nutrient sequestration in the conversion from grassland or savanna to woodland (Johnston *et al.*, 1996; Scholes and van der Merwe, 1996; Scholes and Bailey, 1996; San José *et al.*, 1998), these lands may also have "carbon credit" value to society (e.g., Glenn *et al.*, 1992). Government or industry subsidies and payments for management practices that promote or maintain woody plant cover on rangelands would stand in sharp contrast to past rangeland management practices that have sought to eliminate or reduce woody vegetation cover using costly and often short-lived chemical or mechanical treatments that may not produce desired results (Belsky, 1996) and may convert landscapes from sinks to sources of greenhouse gases (De Castro and Kauffman, 1998). Thus, the perspective on woody plants in rangelands may shift from negative (an expensive management problem) to positive (a potential commodity).

From a biogeochemical perspective, potential benefits of C and N sequestration should be weighed against the potentially undesirable absolute increases in NO and NMHC fluxes that may accompany increases in woody plant biomass. The effects of vegetation change on the hydrological cycle must also be considered. The extent to which shifts from herbaceous to woody plant domination might reduce stem flow and groundwater/aquifer recharge remains controversial. In addition, potential increases in ecosystem transpiration associated with woody communities with high LAI and deep root systems might increase atmospheric water vapor and either offset (due to radiative properties of water) or augment (via cloud formation) benefits of C sequestration on greenhouse gas budgets. From a biodiversity perspective, shrubland/woodland communities may be more (La Copita scenario) or less diverse (many *Juniperus* communities)

than grasslands. In either case, the diversity in terms of the kinds of organisms present would be very different. The development of shrublands and woodlands would be at the expense of grassland habitats and the plants and animals characteristic of such habitats. From a socioeconomic perspective, the promotion or maintenance of woody plant biomass on grasslands and savannas would necessitate a radical change in traditional land use and range management perspectives. Revenues generated from marketing of soil and plant carbon stocks could help fund needed restoration efforts and spur economic development in many sectors, but may be socioeconomically disruptive on other fronts (Trexler and Meganck, 1993). Thus, there are many important issues to be resolved when evaluating the merits of C and N sequestration associated with vegetation change.

5. Implications for Ecosystem and Natural Resources Management

Woody plant encroachment has been and continues to be a major problem in grasslands and savannas worldwide (e.g., Grossman and Gandar, 1989). Because of its direct effects on livestock production,




FIGURE 11 Feedbacks between climate and atmospheric processes, ecosystem structure and function, and human activities. Dashed lines depict traditional research and public awareness; solid lines denote areas requiring increased research emphasis. Understanding global change will ultimately hinge on understanding how socioeconomics, policy, and government subsidies influence human activities and land use (see Figs. 12 and 13).



FIGURE 12 LANDSAT image (1978) of the Canada (Saskatchewan/Alberta)–U.S. (Montana) border in the vicinity of the Milk River (from Knight, 1991). Subsequent to the drought and Dust Bowl of the 1930s, farmlands in Canada were repossessed by provincial or federal governments, withdrawn from cultivation, and underwent secondary succession. Intensive agriculture was maintained in the United States via elaborate farm subsidy programs. Striking contrasts in regional land cover were thus a direct result of changes in government policy. See also color insert.

encroachment of woody vegetation into grasslands has been one of the most important problems facing the ranching industry in the western United States and graziers and pastoralists in arid/semiarid regions throughout the world. This structural change in vegetation also has profound effects on the functional properties of ecosystems. Since woody plant encroachment into grasslands is occurring over large areas worldwide, ecosystem-level changes in nutrient pool sizes and fluxes will likely have important ramifications for regional and global biogeochemistry and climate. Thus, the replacement of grassland/savanna ecosystems by woodlands should be viewed not only as a local problem with economic impacts on livestock husbandry, but also in the context of longer-term, regional impacts on biogeochemistry and climate that will influence future land use options in arid and semiarid ecosystems worldwide.

Since the dawn of time, humans have been cognizant of the direct (e.g., catastrophic floods, wind storms, hail) and indirect (e.g., drought effects on food availability) effects of climate on their well-being. Ecosystem science in the 1960s and 70s focused on climatic and abiotic controls over ecosystem structure and function. During this same period, it became increasingly clear that human activities were directly responsible for significant changes in atmospheric chemistry which could feed back to affect ecosystem processes (e.g., acid rain) and human health (e.g., smog). While ecologists and natural resource managers have long been concerned with impacts of humans on ecosystems, we have only recently begun to assess how alterations of ecosystem structure and function might induce changes in climate and atmospheric chemistry, as shown in Figure 11 (Graetz, 1991; Bryant *et al.*, 1990; Pielke *et al.*, 1993, 1998). The case study presented here explicitly documents how human activities (specifically alteration of grazing and fire regimes) have modified the structure and function of a subtropical savanna grassland system in ways that may have significant impacts on climate and atmospheric processes. Anticipating future changes will largely depend on anticipating how human populations and land use will change. Land-use practices will be governed largely by socioeconomic conditions mediated by government policy and subsidies (Figs. 12 (see also color insert) and 13). Thus, the human dimension of global change is paramount and ecologists are challenged to interface ecosystem science with social science (Turner *et al.*, 1990; Walker, 1993a, b; Walker and Steffen, 1993; Vitousek, 1994).

Changes in tree-shrub-grass ratios in drylands have policy implications for federal agencies grappling with designing and implementing carbon sequestration programs. The success of such endeavors will hinge on the ability to quantitatively monitor and inventory "carbon credits" associated with various land management practices. In arid and semiarid ecosystems, this means tracking changes in woody versus herbaceous cover and understanding how shifts between these growth forms influences aboveground and belowground C and N pools and fluxes. However, landscape-scale and regional quantification of grass-woody plant mass is challenging, because woody encroachment occurs relatively slowly (decadal time scales) in a nonlinear fashion, across large and often remote areas, and in a heterogeneous manner determined by topographic constraints, climate, land-use, and disturbance regimes. In addition, reductions in woody biomass also occur in




FIGURE 13 Differences in biophysical properties associated with contrasting land management policies in adjacent portions of the United States and Mexico (adapted from Bryant *et al.*, 1990). Relaxation of grazing and range improvement programs occurred in the United States subsequent to the implementation of the Taylor Grazing Act in 1934. Changes in landscape cover resulting from a change in federal policy has had biophysical consequences.

drylands where trees and shrubs are cleared using fire, herbicides, or mechanical means (e.g., roller chopping, chaining). As a result, landscapes within a region may be a mosaic of variable-strength C and N sources and sinks. At present, we lack comprehensive information on the rate of change, areal extent, and pattern of woody plant abundance in the world's drylands. Hence, it is difficult to objectively assess the role of savannas in regional/global C and N cycling. Recent advances in remote sensing show promise for quantifying changes in grass and woody plant biomass in drylands (Asner *et al.*, 1998b). These remote sensing tools, when used in conjunction with simulation modeling (Asner *et al.*, 1998a), will potentially enable functional monitoring of land-use impacts on regional biogeochemistry in savanna regions.

6. Summary

Woody plant encroachment has been widespread in grassland and savanna ecosystems over the past century. This phenomenon jeopardizes grassland biodiversity and threatens the sustainability of pastoral, subsistence and commercial livestock grazing. As such, it may adversely impact ~20% of the world's population. Although woody plant expansion has long been a concern of land managers

in grassland and savanna regions, most research on this issue has focused on woody plant effects on grass production and the development methods to limit or reduce the abundance of trees and shrubs. Little is known of the rates and dynamics of the phenomenon or its impact on fundamental ecological processes related to energy flow and nutrient cycling. Grassland/savanna systems account for 30–35% of global terrestrial net primary production. Hence, when woody species increase in abundance and transform grasslands and savannas into shrublands and woodlands, the potential to alter land surface–atmosphere interactions and C and N sequestration and cycling at regional and global scales may be significant.

The La Copita case study documents the rate and magnitude of change in ecosystem biogeochemistry that can occur when a subtropical dryland landscape is transformed from savanna grassland to woodland. Linked succession–biogeochemistry models, confirmed with historical aerial photography and ground measurements, indicate that soil and plant C mass has increased 10% and 10-fold, respectively, with succession from presettlement savanna grassland to present-day savanna woodland. Ecosystem C storage will continue to increase as present-day woody vegetation communities mature and expand into remaining herbaceous areas. Accumulation of ecosystem C mass was accompanied by increases in soil N pools. Fluctuations in monthly woody plant root biomass in upper soil horizons exceeded foliar litter inputs by one to two orders of magnitude, suggesting that belowground inputs of organic matter drive changes in soil physical and chemical properties subsequent to woody plant establishment in grasslands. The deep root systems of woody plants have also increased C mass throughout the soil profile relative to that of grasslands. Increases in C and N pools have occurred in spite of increases in N mineralization, NO flux, soil respiration, and nonmethane hydrocarbon emissions.

These results are of potential global significance, given that large areas of Africa, South America, North America, and Australia have been undergoing similar land cover changes over the past century. The demonstrated capacity for carbon sequestration in this semiarid system suggests a need to reevaluate traditional perspectives on woody plants in rangelands as governments and industries seek ways to mitigate greenhouse gas emissions. However, sequestration of C by woody plants in drylands may come at the expense of elevated NO_x, NMHC, and groundwater fluxes. Regional assessments of the potential consequences of global change are hampered by a lack of quantitative information on the geographic balance between woody plant expansion and clearing in the world's extensive and often remote drylands. Recent developments in linked remote-sensing ecosystem modeling approaches show promise for alleviating these monitoring constraints.

Acknowledgments

We thank Chad McMurtry for assisting with figures and with computation of field estimates in Figure 7. Xiaolian Ren assisted with the image processing and GIS used to generate the data in Figures 4 and 5. This research was supported by grants from

NASA-EOS (NAGW-2662), NASA-LCLUC (NAG5-6134), the NASA Graduate Student Research Program, DOE-NIGEC (DE-F03-90ER61010), USDA-NRICGP (96-00842), and the Texas ATRP (999902126).

References

- Abrams, M. D. (1986). Historical development of gallery forests in north-east Kansas. *Vegetatio* 65, 29–37.
- Acoccks, J. P. H. (1964). Karroo vegetation in relation to the development of deserts. In "Ecological Studies of Southern Africa" (D. H. S. Davis, Ed.), pp. 100–112. Dr. Junk Publishers, The Hague.
- Adamoli, J., Sennhauser, E., Acero, J. M., and Rescia, A. (1990). Stress and disturbance: Vegetation dynamics in the dry Chaco region of Argentina. *J. Biogeogr.* 17, 491–500.
- Ambrose, S. H., and Sikes, N. E. (1991). Soil carbon isotope evidence for Holocene habitat change in the Kenya Rift Valley. *Science* 253, 1402–1405.
- Anderson, J. E., and Holte, K. E. (1981). Vegetation development over 25 years without grazing on sagebrush-dominated rangeland in southeastern Idaho. *J. Range Manage.* 31, 25–29.
- Anderson, R. C., Erlich, J. S., and Baskin, J. M. (Eds.). (1999). "Savanna, barren and rock outcrop plant communities of North America." Cambridge University Press, New York.
- Ansley, J. A., Pinchak, W. E., and Ueckert, D. N. (1995). Changes in red-berry juniper distribution in northwest Texas (1948–1982). *Rangelands* 17, 49–53.
- Archer, S. (1989). Have southern Texas savannas been converted to woodlands in recent history? *Am. Nat.* 134, 545–561.
- Archer, S. (1994). Woody plant encroachment into southwestern grasslands and savannas: rates, patterns and proximate causes. In "Ecological Implications of Livestock Herbivory in the West" (M. Vavra, W. Laycock, and R. Pieper, Eds.), pp. 13–68. Society for Range Management, Denver, Colorado.
- Archer, S. (1995a). Herbivore mediation of grass-woody plant interactions. *Trop. Grassl.* 29, 218–235.
- Archer, S. (1995b). Tree-grass dynamics in a *Prosopis*-thornscrub savanna parkland: reconstructing the past and predicting the future. *Ecoscience* 2, 83–99.
- Archer, S. (1996). Assessing and interpreting grass-woody plant dynamics. In "The Ecology and Management of Grazing Systems" (J. Hodgson and A. Illius, Eds.), pp. 101–134. CAB International, Wallingford, Oxon, United Kingdom.
- Archer, S., Schimel, D. S., and Holland, E. A. (1995). Mechanisms of shrubland expansion: land use, climate or CO₂? *Climatic Change* 29, 91–99.
- Archer, S., Scifres, C. J., Bassham, C. R., and Maggio, R. (1988). Autogenic succession in a subtropical savanna: conversion of grassland to thorn woodland. *Ecol. Monogr.* 58, 111–127.
- Archer, S. and Stokes, C. J. (2000). Stress, disturbance and change in rangeland ecosystems. In "Rangeland Desertification" (O. Arnalds and S. Archer, Eds.), pp. 17–38. Advances in Vegetation Science Vol. 19, Kluwer Publishing Company, Dordrecht, Netherlands.
- Archibold, O. W., and Wilson, M. R. (1980). The natural vegetation of Saskatchewan prior to agricultural settlement. *Can. J. Bot.* 58, 2031–2042.
- Arnalds, O. and Archer, S. (Eds.). (2000). "Rangeland Desertification." Advances in Vegetation Science Vol. 19. Kluwer Publishing Company, Dordrecht, Netherlands.

- Arno, S. F. and Gruell, G. E. (1986). Douglas fir encroachment into mountain grasslands in southwestern Montana. *J. Range Manage.* 39, 272–276.
- Arno, S. F., Harrington, M. G., Fiedler, C. E., and Carlson, C. E. (1995). Restoring fire-dependent ponderosa pine forests in western Montana. *Restoration and Management Notes* 13, 32–36.
- Arnold, J. F. (1950). Changes in ponderosa pine bunchgrass ranges in northern Arizona resulting from pine regeneration and grazing. *J. Forestry* 48, 118–126.
- Asner, G. P., Bateson, C. A., Privette, J. L., El Saleous, N., and Wessman, C. A. (1998a). Estimating vegetation structural effects on carbon uptake using satellite data fusion and inverse modeling. *J. Geophys. Res.* 103, 28,839–28,853.
- Asner, G. P., Wessman, C. A., and Schimel, D. S. (1998b). Heterogeneity of savanna canopy structure and function from imaging spectrometry and inverse modeling. *Ecol. Appl.* 8, 1022–1036.
- Backéus, I. (1992). Distribution and vegetation dynamics of humid savannas in Africa and Asia. *J. Veg. Sci.* 3, 345–356.
- Bahre, C. J. (1991). "A Legacy of Change: Historic Human Impact on Vegetation of the Arizona Borderlands." University of Arizona Press, Tucson.
- Bahre, C. J. and Shelton, M. L. (1993). Historic vegetation change, mesquite increases, and climate in southeastern Arizona. *J. Biogeogr.* 20, 489–504.
- Baker, W. L. and Weisberg, P. J. (1997). Using GIS to model tree population parameters in the Rocky Mountain National Park forest-tundra ecotone. *J. Biogeogr.* 24, 513–526.
- Barnes, P. W. and Archer, S. R. (1996). Influence of an overstorey tree (*Prosopis glandulosa*) on associated shrubs in a savanna parkland: implications for patch dynamics. *Oecologia* 105, 493–500.
- Barnes, P. W. and Archer, S. R. (1998). Tree-shrub interactions in a subtropical savanna parkland: Competition or facilitation? *J. Veg. Sci.* 10, 525–536.
- Bauer, A., Cole, C. V., and Black, A. L. (1987). Soil property comparisons in virgin grasslands between grazed and nongrazed management systems. *Soil Sci. Soc. Am. J.* 51, 176–182.
- Belsky, A. J. (1996). Western juniper expansion: is it a threat to arid northwestern ecosystems? *J. Range Manage.* 49, 53–59.
- Ben-Shahar, R. (1991). Successional patterns of woody plants in catchment areas in a semi-arid region. *Vegetatio* 93, 19–27.
- Bernhard-Reversat, F. (1982). Biogeochemical cycle of nitrogen in a semi-arid savanna. *Oikos* 38, 321–332.
- Bews, J. W. (1917). Plant succession in the thorn veld. *S. Afr. J. Sci.* 4, 153–172.
- Binggeli, P. (1996). A taxonomic, biogeographical and ecological overview of invasive woody plants. *J. Veg. Sci.* 7, 121–124.
- Blackburn, W. H. and Tueller, P. T. (1970). Pinyon and juniper invasion in black sagebrush communities in east-central Nevada. *Ecology* 51, 841–848.
- Blair, W. F. (1950). The biotic provinces of Texas. *Texas J. Sci.* 2, 93–117.
- Bock, J. H. and Bock, C. E. (1984). Effect of fires on woody vegetation in the pine-grassland ecotone of the southern Black Hills. *Am. Midl. Nat.* 112, 35–42.
- Bogusch, E. R. (1952). Brush invasion of the Rio Grande Plains of Texas. *Texas J. Sci.* 4, 85–91.
- Bonan, G. B. (1997). Effects of land use on the climate of the United States. *Climatic Change* 37, 449–486.
- Bond, W. J., Stock, W. D., and Hoffman, M. T. (1994). Has the Karoo spread? A test for desertification using carbon isotopes from soils. *S. Afr. J. Sci.* 90, 391–397.
- Booth, C. A. and Barker, P. J. (1981). Shrub invasion on sandplain country west of Wanaaring, New South Wales. *Soil Conservation Service of New South Wales* 37, 65–70.
- Bossard, C. C. (1991). The role of habitat disturbance, seed predation and ant dispersal on establishment of the exotic shrub *Cytisus scoparius* in California. *Am. Midl. Nat.* 126, 1–13.
- Bossard, C. C. and Rejmanek, M. (1994). Herbivory, growth, seed production, and resprouting of an exotic invasive shrub. *Biol. Conserv.* 67, 193–200.
- Boutton, T., Archer, S., and Midwood, A. (1999). Stable isotopes in ecosystem science: structure, function and dynamics of a subtropical savanna. *Rapid Commun. Mass Spectrom.* 13, 1–15.
- Boutton, T. W., Archer, S. R., Midwood, A. J., Zitzer, S. F., and Bol, R. (1998). $\delta^{13}\text{C}$ values of soil organic carbon and their use in documenting vegetation change in a subtropical savanna ecosystem. *Geoderma* 82, 5–41.
- Bowman, D. M. J. S. and Panton, W. J. (1995). Munmarlary revisited: response of a north Australian *Eucalyptus tetrodonta* savanna protected from fire for 20 years. *Aust. J. Ecol.* 20, 526–531.
- Bragg, T. B. and Hulbert, L. C. (1976). Woody plant invasion of unburned Kansas bluestem prairie. *J. Range Manage.* 29, 19–24.
- Branscomb, B. L. (1958). Shrub invasion of a southern New Mexico desert grassland range. *J. Range Manage.* 11, 129–132.
- Bray, W. L. (1901). The ecological relations of the vegetation of western Texas. *Bot. Gaz.* 32, 99–123; 195–217; 262–291.
- Bren, L. J. (1992). Tree invasion of an intermittent wetland in relation to changes in the flooding frequency of the Murray River, Australia. *Aust. J. Ecol.* 17, 395–408.
- Biggs, J. M. and Gibson, D. G. (1992). Effects of burning on tree spatial patterns in a tallgrass prairie landscape. *Bull. Torr. Bot. Club* 119, 300–307.
- Brown, A. L. (1950). Shrub invasions of southern Arizona desert grasslands. *J. Range Manage.* 3, 172–177.
- Brown, D. (1994). The impact of species introduced to control tree invasion on the vegetation of an electrical utility right-of-way. *Can. J. Bot.* 73, 1217–1228.
- Brown, J. H., Valone, T. J., and Curtin, C. G. (1997). Reorganization of an arid ecosystem in response to recent climate change. *Proc. Natl. Acad. Sci.* 94, 9729–9733.
- Brown, J. R. and Archer, S. (1987). Woody plant seed dispersal and gap formation in a North American subtropical savanna woodland: the role of domestic herbivores. *Vegetatio* 73, 73–80.
- Brown, J. R. and Archer, S. (1990). Water relations of a perennial grass and seedling versus adult woody plants in a subtropical savanna, Texas. *Oikos* 57, 366–374.
- Brown, J. R. and Archer, S. (1999). Shrub invasion of grassland: recruitment is continuous and not regulated by herbaceous biomass or density. *Ecology* 80, 2385–2396.
- Brown, J. R. and Carter, J. (1998). Spatial and temporal patterns of exotic shrub invasion in an Australian tropical grassland. *Landscape Ecol.* 13, 93–103.
- Bruce, K., Cameron, G., and Harcombe, P. (1995). Initiation of a new woodland type on the Texas coastal prairie by the chinese tallow tree (*Sapium sebiferum* (L) Roxb). *Bull. Torr. Bot. Club* 122, 215–225.
- Bryant, N. A., Johnson, L. F., Brazel, A. J., Balling, R. C., Hutchinson, C. F., and Beck, L. R. (1990). Measuring the effect of overgrazing in the Sonoran Desert. *Climatic Change* 17, 243–264.
- Bücher, E. H. (1982). Chaco and Caatinga-South American arid savannas, woodlands, and thickets. In "Ecology of Tropical Savannas" (B. J.

- Huntley and B. H. Walker, Eds.), pp. 48–79. Springer-Verlag, New York.
- Bücher, E. H. (1987). Herbivory in arid and semi-arid regions of Argentina. *Revista Chilena de Historia Natural* 60, 265–273.
- Buffington, L. D. and Herbel, C. H. (1965). Vegetational changes on a semidesert grassland range from 1958 to 1963. *Ecol. Monogr.* 35, 139–164.
- Burgess, T. L. (1995). Desert grassland, mixed-shrub savanna, shrub-steppe, or semi-desert scrub? The dilemma of coexisting growth forms. In "The Desert Grasslands" (M. P. McClaran, and T. R. Van Devender, Eds.), pp. 31–67. University of Arizona Press, Tucson.
- Burkhardt, J. and Tisdale, E. W. (1976). Causes of juniper invasion in southwestern Idaho. *Ecology* 57, 472–484.
- Burrows, W. H., Beale, I. F., Silcock, R. G., and Pressland, A. J. (1985). Prediction of tree and shrub population changes in a semi-arid woodland. In "Ecology and Management of the World's Savannas" (J. C. Tothill and J. J. Mott, Eds.), pp. 207–211. Australian Academy of Science, Canberra.
- Burrows, W. H., Carter, J. O., Scanlan, J. C., and Anderson, E. R. (1990). Management of savannas for livestock production in north-east Australia: contrasts across the tree-grass continuum. *J. Biogeogr.* 17, 503–512.
- Burrows, W. H., Compton, J. F., and Hoffmann, M. B. (1998). Vegetation thickening and carbon sinks in the grazed woodlands of north-east Australia. In "Proceedings, Australian Forest Growers Conference." pp. 305–316. Lismore.
- Callaway, R. M. and Davis, F. W. (1993). Vegetation dynamics, fire, and the physical environment in coastal Central California. *Ecology* 74, 1567–1578.
- Canadell, J., Jackson, R. B., Ehleringer, J. R., Mooney, H. A., Sala, O. E., and Schulze, E.-D. (1996). Maximum rooting depth of vegetation types at the global scale. *Oecologia* 108, 583–595.
- Charley, J. L. and West, N. E. (1975). Plant-induced soil chemical patterns in shrub-dominated semi-desert ecosystems of Utah. *J. Ecol.* 63, 945–964.
- Ciais, P., Tans, P. P., Trolier, M., White, J. W. C., and Francey, R. J. (1995). A large northern hemisphere terrestrial CO₂ sink indicated by the ¹³C/¹²C ratio of atmospheric CO₂. *Science* 269, 1098–1102.
- Cline-Cole, R. A., Main, H. A. C., and Nichol, J. E. (1990). On fuelwood consumption, population dynamics and deforestation in Africa. *World Dev.* 18, 513–527.
- Cole, J. K., Martin, R. E., Holland, E. A., Archer, S. R., Hibbard, K., and Scholes, M. (1996). Nitric oxide fluxes from a subtropical savanna. La Copita Research Area 1996 Consolidated Progress Report CPR-5047, Texas Agricultural Experiment Station.
- Coley, P. D., Bryant, J. P., and Chapin, F. S. III. (1985). Resource availability and plant antiherbivore defense. *Science* 230, 895–899.
- Connin, S. L., Virginia, R. A., and Chamberlain, C. P. (1997). Carbon isotopes reveal soil organic matter dynamics following arid land shrub expansion. *Oecologia* 110, 374–386.
- Cook, G. D., Setterfield, S. A., and Maddison, J. P. (1996). Shrub invasion of a tropical wetland: implications for weed management. *Ecol. Appl.* 6, 531–537.
- Cooper, C. F. (1960). Changes in vegetation, structure, growth of southwestern pine forests since white settlement. *Ecol. Monogr.* 30, 129–164.
- Covington, W. W. and Moore, M. M. (1994). Southwestern ponderosa forest structure: changes since Euro-American settlement. *J. Forestry* 92, 39–47.
- Crutzen, P. J. and Andreae, M. O. (1990). Biomass burning in the tropics: impacts on atmospheric chemistry and biogeochemical cycles. *Science* 250, 1669–1678.
- Crutzen, P. J. and Goldammer, J. G. (1993). "Fire in the Environment: The Ecological, Atmospheric, and Climatic Importance of Vegetation Fires." John Wiley, New York.
- Cunningham, G. M. and Walker, P. J. (1973). Growth and survival of mulga (*Acacia aneura* F. Muell. ex Benth) in western New South Wales. *Trop. Grassl.* 7, 69–77.
- D'Antonio, C. M. and Vitousek, P. M. (1992). Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annu. Rev. Ecol. System.* 23, 63–87.
- De Castro, E. A. and Kauffman, J. B. (1998). Ecosystem structure in the Brazilian Cerrado: a vegetation gradient of aboveground biomass, root mass and consumption by fire. *J. Trop. Ecol.* 14, 263–283.
- De Steven, D. (1991). Experiments on mechanisms of tree establishment in old-field succession: seeding survival and growth. *Ecology* 72, 1076–1088.
- Dick-Peddie, W. A. (1993). "New Mexico vegetation: past, present, and future." University of New Mexico Press, Albuquerque, New Mexico.
- Distel, R. A. and Boo, R. M. (1996). Vegetation states and transitions in temperate semi-arid rangelands of Argentina. In "Proceedings, Vth International Rangeland Congress, Vol. II" (N. E. West, Ed.), pp. 117–118. Society for Range Management, Salt Lake City, Utah.
- Dussart, E., Lerner, P., and Peinetti, R. (1998). Long-term dynamics of two populations of *Prosopis caldenia* Burkart. *J. Range Manage.* 51, 685–691.
- Eissenstat, D. M. and Yanai, R. D. (1997). The ecology of root life spans. *Adv. Ecol. Res.* 27, 1–60.
- Ellis, D. and Schuster, J. L. (1968). Juniper age and distribution on an isolated butte in Garza County. *Southwest. Nat.* 13, 343–348.
- Engle, D. M., Bidwell, T. G., and Moseley, M. E. (1996). Invasion of Oklahoma rangelands and forests by eastern redcedar and ashe juniper. Circular E-947, Coop. Extension Service, Oklahoma State University, Stillwater, Oklahoma.
- Fehsenfeld, F., Calvert, J., Fall, R., Goldan, P., Guenther, A. B., Hewitt, C. N., Lamb, B., Liu, S., Trainer, M., Westberg, H., and Zimmerman, P. (1992). Emissions of volatile organic compounds from vegetation and the implications for atmospheric chemistry. *Global Biogeochem. Cycles* 6, 389–430.
- Field, C. B., Behrenfeld, M. J., Randerson, J. T., and Falkowski, P. (1998). Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* 281, 237–240.
- Fisher, C. E. (1950). The mesquite problem in the Southwest. *J. Range Manage.* 3, 60–70.
- Fisher, C. E. (1977). Mesquite and modern man in southwestern North America. In "Mesquite: Its Biology in Two Desert Ecosystems" (B. B. Simpson, Ed.), pp. 177–188. US/IBP Synthesis Series Vol. 4, Dowden, Hutchinson & Ross, New York.
- Flinn, R. C., Scifres, C. J., and Archer, S. R. (1992). Variation in basal sprouting in co-occurring shrubs: implications for stand dynamics. *J. Veg. Sci.* 3, 125–128.
- Foster, J. H. (1917). The spread of timbered areas in central Texas. *J. Forestry* 15, 442–445.
- Frank, A. B., Tanaka, D. L., Hofmann, L., and Follett, R. F. (1995). Soil carbon and nitrogen of northern Great Plains grasslands as influenced by long-term grazing. *J. Range Manage.* 48, 470–474.
- Friedel, M. H. (1987). A preliminary investigation of woody plant increase in the western Transvaal and implications for veld assessment. *J. Grassl. Soc. S. Afr.* 4, 25–30.
- Gadgil, M. and Meher-Homji, V. M. (1985). Land use and productive potential of Indian savannas. In "Ecology and Management of the World's


- Savannas" (J. C. Tothill and J. J. Mott, Eds.), pp. 107–113. Australian Academy of Science, Canberra.
- Gardener, G. J., McIvor, J. G., and Williams, J. (1990). Dry tropical rangelands: solving one problem and creating another. In "Australian Ecosystems: 200 Years of Utilization, Degradation and Reconstruction" (D. A. Saunders, A. J. M. Hopkins, and R. A. How, Eds.), pp. 279–286. Ecological Society of Australia, Geraldton, Western Australia.
- Gardiner, C. P. and Gardiner, S. P. (1996). The dissemination of chinese apple (*Ziziphus mauritiana*): a woody weed of the tropical subhumid savanna and urban fringe of north Queensland. *Trop. Grassl.* 30, 174–174.
- Gibbens, R. P., Beck, R. F., McNeely, R. P., and Herbel, C. H. (1992). Recent rates of mesquite establishment on the northern Chihuahuan Desert. *J. Range Manage.* 45, 585–588.
- Gill, R. A. and Burke, I. C. (1999). Ecosystem consequences of plant life form changes at three sites in the semiarid United States. *Oecologia* 122, 551–563.
- Glendening, G. E. (1952). Some quantitative data on the increase of mesquite and cactus on a desert grassland range in southern Arizona. *Ecology* 33, 319–328.
- Glendening, G. E. and Paulsen, H. A. (1955). Reproduction and establishment of velvet mesquite as related to invasion of semi-desert grasslands. Technical Bulletin 1127, U.S. Department of Agriculture, Washington, DC.
- Glenn, E. P., Pitelka, L. F., and Olsen, M. W. (1992). The use of halophytes to sequester carbon. *Water, Air and Soil Pollution* 64, 251–263.
- Graetz, R. D. (1991). The nature and significance of the feedback of changes in terrestrial vegetation on global atmospheric and climatic change. *Climatic Change* 18, 147–173.
- Grice, A. C. (1996). Seed production, dispersal and germination in *Cryptostegia grandiflora* and *Ziziphus mauritiana*, two invasive shrubs in tropical woodlands of northern Australia. *Aust. J. Ecol.* 21, 324–331.
- Grice, A. C. (1997). Post-fire regrowth and survival of the invasive tropical shrubs *Cryptostegia grandiflora* and *Ziziphus mauritiana*. *Aust. J. Ecol.* 22, 49–55.
- Grimm, E. C. (1983). Chronology and dynamics of vegetation change on the prairie-woodland region of southern Minnesota, USA. *New Phytol.* 93, 311–350.
- Grossman, D. and Gandar, M. V. (1989). Land transformation in South African savannah regions. *S. Afr. Geogr. J.* 71, 38–45.
- Grover, H. D. and Musick, H. B. (1990). Shrubland encroachment in southern New Mexico, USA: an analysis of desertification processes in the American southwest. *Climatic Change* 17, 305–330.
- Gruell, G. E. (1983). Fire and vegetative trends in the northern Rockies: interpretations from 1871–1982 photographs. INT-158, Ogden, Utah, USDA Forest Service Intermountain Research Station General Technical Report.
- Guenther, A. (1997). Seasonal and spatial variations in natural volatile organic compound emissions. *Ecol. Appl.* 7, 34–45.
- Guenther, A., Archer, S., Greenberg, J., Harley, P., Helmig, D., Klinger, L., Vierling, L., Wildermuth, M., Zimmerman, P., and Zitzer, S. (1999). Biogenic hydrocarbon emissions and landcover/climate change in a subtropical savanna. *Phy. Chem. Earth (B)* 24, 659–667.
- Guenther, A., Hewitt, C., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P. (1995). A global model of natural volatile organic compound emissions. *J. Geophys. Res.* 100, 8873–8892.
- Hall, D. O., Ojima, D. S., Parton, W. J., and Scurlock, J. M. O. (1995). Response of temperate and tropical grasslands to CO₂ and climate change. *J. Biogeogr.* 22, 537–547.
- Handley, L. L. and Scrimgeour, C. M. (1997). Terrestrial plant ecology and ¹⁵N abundance: the present limits to interpretation for uncultivated systems with original data from a Scottish old field. *Adv. Ecol. Res.* 27, 133–213.
- Hao, W. M., Liu, M. H., and Crutzen, P. J. (1990). Estimates of annual and regional releases of CO₂ and other trace gases to the atmosphere from fires in the tropics, based on the FAO statistics for the period 1975–1980. In "Fire in the Tropical Biota" (J. G. Goldammer, Ed.), pp. 440–462. Springer-Verlag, New York.
- Harrington, G. N. and Hodgkinson, K. C. (1986). Shrub-grass dynamics in mulga communities of eastern Australia. In "Rangelands: A Resource Under Siege" (P. J. Joss, P. W. Lynch, and O. B. Williams, Eds.), pp. 26–28. Australian Academy of Science, Canberra.
- Harrington, G. N., Oxley, R. E., and Tongway, D. J. (1979). The effects of European settlement and domestic livestock on the biological system in poplar box (*Eucalyptus populnea*) lands. *Aust. Rangel. J.* 1, 271–279.
- Hart, R. H. and Laycock, W. A. (1996). Repeat photography on range and forest lands in the western United States. *J. Range Manage.* 49, 60–67.
- Hastings, J. R. and Turner, R. L. (1965). "The Changing Mile: An Ecological Study of Vegetation Change with Time in the Lower Mile of an Arid and Semi-Arid Region," University of Arizona Press, Tucson.
- Hendricks, J. J., Nadelhoffer, K. J., and Aber, J. D. (1997). A N-15 tracer technique for assessing fine root production and mortality. *Oecologia* 112, 300–304.
- Hennessy, J. T., Gibbens, R. P., and Tromble, J. M. A. C. M. (1983). Vegetation changes from 1935 to 1980 in mesquite dunelands and former grasslands of southern New Mexico. *J. Range Manage.* 36, 370–374.
- Hibbard, K. A. (1995). Landscape patterns of carbon and nitrogen dynamics in a subtropical savanna: observations and models. Ph.D. dissertation, Texas A&M University, College Station.
- Hobbs, R. J. and Mooney, H. A. (1986). Community changes following shrub invasion of grassland. *Oecologia* 70, 508–513.
- Höchberg, M. E., Menaut, J.-C., and Gignoux, J. (1994). The influences of tree biology and fire in the spatial structure of a West African savannah. *J. Ecol.* 82, 217–226.
- Hodgkin, S. E. (1984). Shrub encroachment and its effects on soil fertility on Newborough Warren, Anglesey, Wales. *Biol. Conserv.* 29, 99–119.
- Holland, E. A., Dentener, F. J., Braswell, B. H., and Sulzman, J. M. (1999). Contemporary and pre-industrial global reactive nitrogen budgets. *Biogeochemistry* 46, 7–43.
- Holmes, P. M. and Cowling, R. M. (1997). The effects of invasion by *Acacia saligna* on the guild structure and regeneration capabilities of South African fynbos shrublands. *J. Appl. Ecol.* 34, 317–332.
- Hopkins, B. (1983). Successional processes. In "Tropical Savannas" (F. Bourliere, Ed.), pp. 605–616. Ecosystems of the World Vol. 13, Elsevier, Amsterdam.
- Houghton, R. A., Boone, R. D., Fruci, J. R., Hobbie, J. E., Melillo, J. M., Palm, C. A., Peterson, B. J., Shaver, G. R., Woodwell, G. M., Moore, B., Skoles, D. L., and Myers, N. (1987). The flux of carbon from terrestrial ecosystems to the atmosphere in 1980 due to changes in land use: geographic distribution of the global flux. *Tellus* 39B, 122–139.
- Humphrey, R. R. (1953). The desert grassland, past and present. *J. Range Manage.* 6, 159–164.
- Humphrey, R. R. (1958). The desert grasslands: a history of vegetation change and an analysis of causes. *Bot. Rev.* 24, 193–252.

- Humphrey, R. R. (1987). "90 Years and 535 Miles: Vegetation Changes Along the Mexican Border." University of New Mexico Press, Albuquerque.
- Humphrey, R. R. and Mehrhoff, L. A. (1958). Vegetation change on a southern Arizona grassland range. *Ecology* 39, 720–726.
- Idso, S. B. (1995). "CO₂ and the Biosphere: the Incredible Legacy of the Industrial Revolution." Special Publication, Department of Soil, Water & Climate, University of Minnesota, St. Paul.
- Inglis, J. M. (1964). "A History of Vegetation on the Rio Grande Plains." Texas Parks and Wildlife Department, Austin.
- Jackson, R. B., Canadell, J., Ehleringer, J. R., Mooney, H. A., Sala, O. E., and Schulze, E. D. (1996). A global analysis of root distributions for terrestrial biomes. *Oecologia* 108, 389–411.
- Jeltsch, F., Milton, S. J., Dean, W. R. J., and van Rooyen, N. (1997). Simulated pattern formation around artificial waterholes in the semi-arid Kalahari. *J. Veg. Sci.* 8, 177–188.
- Johnsen, T. N. (1962). One-seed juniper invasion of northern Arizona grasslands. *Ecol. Monogr.* 32, 187–207.
- Johnson, H. B. and Mayeux, H. S. (1990). *Prosopis glandulosa* and the nitrogen balance of rangelands: extent and occurrence of nodulation. *Oecologia* 84, 176–185.
- Johnson, K. L. (1987). Sagebrush over time: a photographic study of rangeland change. In "Biology of *Artemisia* and *Chrysothamnus*" (E. D. McArthur and B. L. Welch, Eds.), pp. 223–252. USDA Forest Service General Technical Report INT-200, Ogden, Utah.
- Johnson, W. C. (1994). Woodland expansion in the Platte River, Nebraska: Patterns and causes. *Ecol. Monogr.* 64, 45–84.
- Johnston, M. C. (1963). Past and present grasslands of southern Texas and northeastern Mexico. *Ecology* 44, 456–466.
- Johnston, M. H., Homann, P. S., Engstrom, J. K., and Grigal, D. F. (1996). Changes in ecosystem carbon storage over 40 years on an old-field/forest landscape in east-central Minnesota. *Forest Ecol. Manage.* 83, 17–26.
- Jones, F. B. (1975). "Flora of the Texas Coastal Bend." Mission Press, Corpus Christi, Texas.
- Kenney, W. R., Bock, J. H., and Bock, C. E. (1986). Responses of the shrub, *Baccharis pteronioides*, to livestock enclosure in southeastern Arizona. *Am. Midl. Nat.* 116, 429–431.
- Klinger, L., Greenberg, J., Guenther, A., Tyndall, G., Zimmerman, P., Bangui, M., Moutsambote, J.-M., and Kenfack, D. (1998). Patterns in volatile organic compound emissions along a savanna-rainforest gradient in Central Africa. *J. Geophys. Res.* 102, 1443–1454.
- Klink, C. A., Moreira, A. G. and Solbrig, O. T. (1993). Ecological impact of agricultural development in the Brazilian *cerrados*. In "The World's Savannas: Economic Driving Forces, Ecological Constraints and Policy Options for Sustainable Land Use" (M. D. Young and O. T. Solbrig, Eds.), pp. 259–282. Man and the Biosphere Vol. 12, Parthenon Publishing Group, Carnforth, United Kingdom.
- Knapp, A. K. and Seastedt, T. R. (1986). Detritus accumulation limits productivity of tallgrass prairie. *Bioscience* 36, 662–668.
- Knapp, P. A. and Soule, P. T. (1996). Vegetation change and the role of atmospheric CO₂ enrichment on a relict site in central Oregon: 1960–1994. *Ann. Assn. Am. Geog.* 86, 387–411.
- Knapp, P. A. and Soule, P. T. (1998). Recent *Juniperus occidentalis* (western juniper) expansion on a protected site in central Oregon. *Global Change Biol.* 4, 347–357.
- Knight, C. L., Briggs, J. M., and Nelis, M. D. (1994). Expansion of gallery forest on Konza Prairie Research Natural Area, Kansas, USA. *Landscape Ecol.* 9, 117–125.
- Knight, D. H. (1991). Congressional incentives for landscape research. *Ecol. Soc. Am. Bull.* 72, 195–203.
- Köchy, M. (1999). Grass-tree interactions in western Canada. Ph.D. Dissertation, University of Regina, Regina, Canada.
- Küchler, A. W. (1964). "The Potential Natural Vegetation of the Conterminous United States." American Geographical Society, New York.
- Le Roux, I. G. (1997). Patterns and rate of woody vegetation cluster development in a semi-arid savanna, KwaZulu-Natal, South Africa. M.S. thesis, Department of Botany, University of Natal.
- Lehman, V. W. (1969). "Forgotten Legions: Sheep in the Rio Grande Plains of Texas." Texas Western University Press, University of Texas at El Paso.
- Leopold, L. B. (1951). Vegetation of southwestern watersheds in the nineteenth century. *Geog. Rev.* 41, 295–316.
- Liao, J., Boutton, T., Hoskisson, A., McCulley, R., and Archer, S. (1999). Woodland development and the N-cycle of a subtropical savanna parkland: insights from $\delta^{15}\text{N}$ of plants and soils. *Abstracts, Ecol. Soc. Am.* 1999, 274.
- Loehle, C., Li, B. L., and Sundell, R. C. (1996). Forest spread and phase transitions at forest-prairie ecotones in Kansas, U.S.A. *Landscape Ecol.* 11, 225–235.
- Lonsdale, W. M. (1993). Rates of spread of an invading species—*Mimosa pigra* in northern Australia. *J. Ecol.* 81, 513–521.
- Madany, M. H. and West, N. E. (1983). Livestock grazing-fire regime interactions within montane forests of Zion National Park, Utah. *Ecology* 64, 661–667.
- Martin, P. H. and Guenther, A. B. (1995). Insights into the dynamics of forest succession and non-methane hydrocarbon trace gas emissions. *J. Biogeogr.* 22, 493–499.
- Martin, S. C. (1975). Ecology and management of southwestern semi-desert grass-shrub ranges: the status of our knowledge. RM-156, Ft. Collins, Colorado, USDA/Forest Service Research Paper.
- Martin, S. C. and Turner, R. M. (1977). Vegetation change in the Sonoran Desert region, Arizona and Sonora. *Arizona Acad. Sci.* 12, 59–69.
- Mast, J. N., Veble, T. T., and Hodgson, M. E. (1997). Tree invasion within a pine/grassland ecotone: an approach with historic aerial photography and GIS modeling. *Forest Ecol. Manage.* 93, 181–194.
- Mast, J. N., Veble, T. T., and Linhart, Y. B. (1998). Disturbance and climatic influences on age structure of ponderosa pine at the pine/grassland ecotone, Colorado Front Range. *J. Biogeogr.* 25, 743–755.
- McBride, J. R. and Heady, H. F. (1968). Invasion of grassland by *Baccharis pilularis* DC. *J. Range Manage.* 21, 106–108.
- McClaran, M. P. and McPherson, G. R. (1995). Can soil organic carbon isotopes be used to describe grass-tree dynamics at a savanna-grassland ecotone and within the savanna? *J. Veg. Sci.* 6, 857–862.
- McCraw, D. J. (1985). Phytogeographic history of *Larrea* in southwestern New Mexico: illustrating the historical expansion of the Chihuahuan Desert. M.A. thesis, University of New Mexico, Albuquerque.
- McCulley, R. (1998). Soil respiration and microbial biomass in a savanna parkland landscape: spatio-temporal variation and environmental controls. M.S. thesis, Texas A&M University, College Station, Texas.
- McKinney, L. B. J. (1996). Forty years of landscape change in Attwater's prairie chicken habitat within the Coastal Prairie of Texas. M.S. thesis, Texas A&M University, College Station, Texas.
- McLendon, T. (1991). Preliminary description of the vegetation of South Texas exclusive of coastal saline zones. *Texas J. Sci.* 43, 13–32.
- McMurtry, C. R. (1997). Gas exchange physiology and water relations of co-occurring woody plant species in a Texas subtropical savanna. M.S. thesis, Southwest Texas State University, San Marcos.
- McPherson, G. (1997). "Ecology and Management of North American Savannas." University of Arizona Press, Tucson, Arizona.
- McPherson, G. R., Boutton, T. W., and Midwood, A. J. (1993). Stable carbon isotope analysis of soil organic matter illustrates vegetation change


- at the grassland/woodland boundary in southeastern Arizona, USA. *Oecologia* 93, 95–101.
- McPherson, G. R., Wright, H. A., and Wester, D. B. (1988). Patterns of shrub invasion in semiarid Texas grasslands. *Am. Midl. Natur.* 120, 391–397.
- Mearns, R. (1995). Institutions and natural resource management: access to and control over woodfuel in East Africa. In "People and Environment in Africa" (T. Binns, Ed.), pp. 103–114. John Wiley, Chichester, United Kingdom.
- Menaut, J. C., Gignoux, J., Prado, C., and Clobert, J. (1990). Tree community dynamics in a humid savanna of the Côte-d'Ivoire: the effects of fire and competition with grass and neighbours. *J. Biogeogr.* 17, 471–481.
- Midwood, A., Boutton, T., Archer, S., and Watts, S. (1998). Water use by woody plants on contrasting soils in a savanna parkland: assessment with $\delta^2\text{H}$ and $\delta^{18}\text{O}$. *Plant Soil* 205, 13–24.
- Milchunas, D. G. and Lauenroth, W. K. (1993). Quantitative effects of grazing on vegetation and soils over a global range of environments. *Ecol. Monogr.* 63, 327–366.
- Milchunas, D. G., Lee, C. A., Lauenroth, W. K., and Coffin, D. P. (1992). A comparison of C-14, Rb-86, and total excavation for determination of root distributions of individual plants. *Plant Soil* 144, 125–132.
- Miller, E. A. and Halpern, C. B. (1998). Effects of environment and grazing disturbance on tree establishment in meadows of the central Cascade Range, Oregon, USA. *J. Veg. Sci.* 9, 265–282.
- Miller, F. H. (1921). Reclamation of grasslands by Utah juniper in the Tusayan National Forest, Arizona. *J. Forestry* 19, 647–657.
- Miller, R. F. and Rose, J. A. (1999). Fire history and western juniper encroachment in sagebrush steppe. *J. Range Manage.* 52, 550–559.
- Miller, R. F. and Rose, J. R. (1995). Historic expansion of *Juniperus occidentalis* southeastern Oregon. *Great Basin Nat.* 55, 37–45.
- Miller, R. F. and Wigand, P. E. (1994). Holocene changes in semiarid pinyon-juniper woodlands. *Bioscience* 44, 465–474.
- Moat, D. A. and Hutchinson, C. F. (Eds.). (1995). "Desertification in Developed Countries." Kluwer Academic Publishers, London.
- Monson, R. K., Guenther, A. B., and Fall, R. (1991). Physiological reality in relation to ecosystem- and global-level estimates of isoprene emission. In "Trace Gas Emissions by Plants" (T. D. Sharkey, E. A. Holland, and H. A. Mooney, Eds.), pp. 185–207. Academic Press, New York.
- Mueller-Dombois, D. and Goldammer, J. B. (1990). Fire in tropical ecosystems and global environmental change: an introduction. In "Fire in the Tropical Biota: Ecosystem Processes and Global Challenges" (J. G. Goldammer, Ed.), pp. 1–10. Springer-Verlag, Berlin.
- Nelson, J. T. and Beres, P. L. (1987). Was it grassland? A look at vegetation in Brewster County, Texas through the eyes of a photographer in 1899. *Texas J. Agri. Nat. Resources* 1, 34–37.
- Noble, J. C. (1997). "The Delicate and Noxious Scrub: Studies on Native Tree and Shrub Proliferation in Semi-Arid Woodlands of Australia." CSIRO Division of Wildlife and Ecology, Canberra.
- Northup, B. K., Zitzer, S. F., Archer, S., and Boutton, T. W. (1996). Direct and indirect estimates of aboveground woody plant biomass. In "La Copita Research Area 1996 Consolidated Progress Report CPR-5047" (J. W. Stuth and S. Dudash, Eds.), pp. 51–53. Texas Agricultural Experiment Station, College Station.
- Norton-Griffiths, M. (1979). The influence of grazing, browsing, and fire on the vegetation dynamics of the Serengeti, Tanzania, Kenya. In "Serengeti: Dynamics of an Ecosystem" (A. R. E. Sinclair and M. Norton-Griffiths, Eds.), pp. 310–352. University of Chicago Press, Chicago, Illinois.
- O'Connor, T. G. and Roux, P. W. (1995). Vegetation changes (1949–71) in a semi-arid, grassy dwarf shrubland in the Karoo, South Africa: influence of rainfall variability and grazing by sheep. *J. Appl. Ecol.* 32, 612–626.
- Ojima, D. S., Bjorn, O. M. D., Glenn, E. P., Owensby, C. E., and Scurlock, J. O. (1993). Assessment of C budget for grasslands and drylands of the world. *Water, Air, and Soil Pollution* 70, 95–109.
- ORNL. (1998). Terrestrial ecosystem responses to global change: a research strategy. Environmental Sciences Division Publication No. 4821, Ecosystems Working Group, Oak Ridge National Laboratories, Oak Ridge, Tennessee.
- Owensby, C. E., Blan, K. R., Eaton, B. J., and Russ, O. G. (1973). Evaluation of eastern redcedar infestations in the northern Kansas Flint Hills. *J. Range Manage.* 26, 256–260.
- Palmer, A. R. and van Rooyen, A. F. (1998). Detecting vegetation change in the southern Kalahari using Landsat TM data. *J. Arid Environ.* 39, 143–153.
- Panetta, F. D. and McKee, J. (1997). Recruitment of the invasive ornamental, *Schinus terebinthifolius*, is dependent upon frugivores. *Aust. J. Ecol.* 22, 432–438.
- Parton, W. J., Ojima, D. S., Cole, C. V., and Schimel, D. S. (1994). A general model for soil organic matter dynamics: sensitivity to litter chemistry, texture, and management. In "Quantitative Modeling of Soil Forming Processes" (R. B. Bryant, Ed.), pp. 147–167. Soil Science Society of America, Madison, Wisconsin.
- Parton, W. J., Schimel, D. S., Cole, C. V., and Ojima, D. S. (1987). Analysis of factors controlling soil organic matter levels in great plains grasslands. *Soil Sci. Soc. Am. J.* 51, 1173–1179.
- Parton, W. J., Scurlock, J. M. O., Ojima, D. S., Gilmanov, T. G., Scholes, R. J., Schimel, D. S., Kirchner, T., Menaut, J.-C., Seastedt, T., Gardia Moya, E., Kamnalrut, A., and Kinyamario, J. I. (1993). Observations and modeling of biomass and soil organic matter dynamics for the grassland biome worldwide. *Global Biogeochem. Cycles* 7, 785–809.
- Parton, W. J., Stewart, J. W. B., and Cole, V. C. (1988). Dynamics of C, N, P and S in grassland soils: a model. *Biogeochemistry* 5, 109–131.
- Pielke, R. A. and Avissar, R. (1990). Influence of landscape structure on local and regional climate. *Landscape Ecol.* 4, 133–155.
- Pielke, R. A., Avissar, R., Raupach, M., Dolman, A. J., Zeng, X. B., and Denning, A. S. (1998). Interactions between the atmosphere and terrestrial ecosystems: influence on weather and climate. *Global Change Biol.* 4, 461–475.
- Pielke, R. A., Schimel, D. S., Kittel, T. G. F., Lee, T. J., and Zeng, X. (1993). Atmosphere-terrestrial ecosystem interactions: implications for coupled modeling. *Ecol. Model.* 67, 5–18.
- Polley, H. W. (1997). Implications of rising atmospheric carbon dioxide concentration for rangelands. *J. Range Manage.* 50, 562–577.
- Polley, H. W., Mayeux, H. S., Johnson, H. B., and Tischler, C. R. (1997). Atmospheric CO₂, soil water, and shrub/grass ratios on rangelands. *J. Range Manage.* 50, 278–284.
- Post, W. M. (1993). Uncertainties in the terrestrial carbon cycle. In "Vegetation Dynamics and Global Change" (A. M. Solomon and H. H. Shugart, Eds.), pp. 116–132. Chapman and Hall, New York.
- Potter, L. D. and Green, D. L. (1964). Ecology of ponderosa pine in western North Dakota. *Ecology* 45, 10–23.
- Progulske, D. R. (1974). "Yellow Ore, Yellow Hair and Yellow Pine: A Photographic Study of a Century of Forest Ecology," South Dakota State University Extension Service, Brookings.
- Raich, J. W. and Schlesinger, W. H. (1992). The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate. *Tellus* 44B, 81–89.
- Ramsay, J. M. and Rose Innes, R. (1963). Some quantitative observations on the effects of fire on the Guinea savanna vegetation of northern Ghana over a period of eleven years. *Afr. Soils* 8, 41–85.
- Rappole, J. H., Russell, C. E., and Fulbright, T. E. (1986). Anthropogenic

- pressures and impacts on marginal, neotropical, semiarid ecosystems: the case of south Texas. *Sci. Total Environ.* 55, 91–99.
- Rapport, D. J. and Whitford, W. G. (1999). How ecosystems respond to stress: common properties of arid and aquatic systems. *Bioscience* 49, 193–203.
- Reichard, S. H. and Hamilton, C. W. (1997). Predicting invasions of woody plants introduced into North America. *Conserv. Biol.* 11, 193–203.
- Reid, R. S. and Ellis, J. E. (1995). Impacts of pastoralists on woodlands in South Turkana, Kenya: livestock-mediated tree recruitment. *Ecol. Appl.* 5, 978–992.
- Reynolds, H. G. and Glendening, G. E. (1949). Merriam kangaroo rat as a factor in mesquite propagation on southern Arizona rangelands. *J. Range Manage.* 2, 193–197.
- Ringose, S., Chanda, R., Musini, N., and Sefe, F. (1996). Environmental change in the mid-Boteti area of north-central Botswana: biophysical processes and human perceptions. *Environ. Manage.* 20, 397–410.
- Robinson, T. W. (1965). Introduction, spread and areal extent of saltcedar (*Tamarix*) in the western States. Professional Paper 491-A, US Geological Survey.
- Rogers, G. F. (1982). "Then and Now: A Photographic History of Vegetation Change in the Central Great Basin Desert." University of Utah Press, Salt Lake City.
- Rummell, R. S. (1951). Some effects of livestock grazing on ponderosa pine forest and range in central Washington. *Ecology* 32, 594–607.
- Sabiiti, E. N. (1988). Fire behaviour and the invasion of *Acacia sieberiana* into savanna grassland openings. *Afr. J. Ecol.* 26, 301–313.
- San José, J. J. and Fariñas, M. R. (1983). Changes in tree density and species composition in a protected *Trachypogon* savanna, Venezuela. *Ecology* 64, 447–453.
- San José, J. J. and Fariñas, M. R. (1991). Temporal changes in the structure of a *Trachypogon* savanna protected for 25 years. *Acta Oecologia* 12, 237–247.
- San José, J. J., Fariñas, M. R., and Rosales, J. (1991). Spatial patterns of trees and structuring factors in a *Trachypogon* savanna of the Orinoco Llanos. *Biotropica* 23, 114–123.
- San José, J. J. and Montes, R. A. (1997). Fire effects on the coexistence of trees and grasses in savannas and the resulting outcome on organic matter budget. *Interciencia* 22, 289–298.
- San José, J. J., Montes, R. A., and Fariñas, M. R. (1998). Carbon stocks and fluxes in a temporal scaling from a savanna to a semi-deciduous forest. *Forest Ecol. Manage.* 105, 251–262.
- Savage, M. and Swetnam, T. W. (1990). Early 19th-century fire decline following sheep pasturing in a Navajo ponderosa pine forest. *Ecology* 71, 2374–2378.
- Scanlan, J. C. (1988). Spatial and temporal vegetation patterns in a subtropical *Prosopis* savanna woodland, Texas. Ph.D. Dissertation, Texas A&M University, College Station.
- Scanlan, J. C. and Archer, S. (1991). Simulated dynamics of succession in a North American subtropical *Prosopis* savanna. *J. Veg. Sci.* 2, 625–634.
- Schlesinger, W. H., Reynolds, J. F., Cunningham, G. L., Huenneke, L. F., Jarrell, W. M., Virginia, R. A., and Whitford, W. G. (1990). Biological feedbacks in global desertification. *Science* 247, 1043–1048.
- Schofield, C. J. and Bucher, E. H. (1986). Industrial contributions to desertification in South America. *Trends Ecol. Evol.* 1, 78–80.
- Scholes, R. C. and Bailey, C. L. (1996). Can savannas help balance the South African greenhouse gas budget. *Afr. J. Sci.* 92, 60–61.
- Scholes, R. J. and Archer, S. R. (1997). Tree-grass interactions in savannas. *Annu. Rev. Ecol. Syst.* 28, 517–544.
- Scholes, R. J. and Hall, D. O. (1996). The carbon budget of tropical savannas, woodlands and grasslands. In "Global Change: Effects on Coniferous Forests and Grasslands" (A. I. Breyer, D. O. Hall, J. M. Melillo, and G. I. Agren, Eds.), pp. 69–100. SCOPE Vol. 56, John Wiley, Chichester, United Kingdom.
- Scholes, R. J. and van der Merwe, M. R. (1996). Sequestration of carbon in savannas and woodlands. *Environ. Professional* 18, 96–103.
- Scholes, R. J. and Walker, B. H. (1993). "An African Savanna: Synthesis of the Nylsvley Study." Cambridge University Press, Cambridge.
- Schüle, W. (1990). Landscapes and climate in prehistory: interaction of wildlife, man, and fire. In "Fire in the Tropical Biota" (J. G. Goldammer, Ed.), pp. 273–315. Springer-Verlag, New York.
- Schwartz, D., de Foresta, H., Mariotti, A., Balesdent, J., Massimba, J. P., and Girardin, C. (1996). Present dynamics of the savanna-forest boundary in the Congolese Mayombe: a pedological, botanical and isotopic (^{13}C and ^{14}C) study. *Oecologia* 106, 516–524.
- Scott, J. D. (1966). Bush encroachment in South Africa. *S. Afr. J. Sci.* 63, 311–314.
- SCS. (1979). Soil Survey of Jim Wells County, Texas. USDA/Soil Conservation Service & Texas Agricultural Experiment Station, College Station, Texas.
- Shantz, H. L. and Turner, B. L. (1958). Photographic documentation of vegetational changes in Africa over a third of a century. College of Agriculture, University of Arizona, Tucson.
- Sinclair, A. R. E. and Fryxell, J. M. (1985). The Sahel of Africa: ecology of a disaster. *Can. J. Zool.* 63, 987–994.
- Skarpe, C. (1990). Structure of the woody vegetation in disturbed and undisturbed arid savanna, Botswana. *Vegetatio* 87, 11–18.
- Skarpe, C. (1991). Spatial patterns and dynamics of woody vegetation in an arid savanna. *J. Veg. Sci.* 2, 565–572.
- Skarpe, C. (1992). Dynamics of savanna ecosystems. *J. Veg. Sci.* 3, 293–300.
- Skovlin, J. M. and Thomas, J. W. (1995). Interpreting long-term trends in Blue Mountain ecosystems from repeat photography. General Technical Report PNW-315, USDA Forest Service Pacific Northwest Research Station.
- Smeins, F. E. (1983). Origin of the brush problem—a geographical and ecological perspective of contemporary distributions. In "Proceedings, Brush Management Symposium" (K. W. McDaniel, Ed.), pp. 5–16. Texas Tech University Press, Lubbock.
- Smeins, F. E., Taylor, T. W., and Merrill, L. B. (1974). Vegetation of a 25-year enclosure on the Edwards Plateau, Texas. *J. Range Manage.* 29, 24–29.
- Smith, A. J. (1975). Invasion and ecesis of bird-disseminated woody plants in a temperate forest sere. *Ecology* 56, 19–34.
- Smith, D. A. and Schmutz, E. M. (1975). Vegetative changes on protected versus grazed desert grassland range in Arizona. *J. Range Manage.* 28, 453–457.
- Snook, E. C. (1985). Distribution of eastern redcedar on Oklahoma rangelands. In "Eastern Redcedar in Oklahoma" (R. F. Wittwer and D. M. Engle, Eds.), pp. 45–52. Publication E-849, Coop. Extension Serv, Division of Agriculture, Oklahoma State University, Stillwater.
- Soule, P. T. and Knapp, P. A. (1999). Western juniper expansion on adjacent disturbed and near-relict sites. *J. Range Manage.* 52, 525–533.
- Steinauer, E. M. and Bragg, T. B. (1987). Ponderosa pine (*Pinus ponderosa*) invasion of Nebraska sandhills prairie. *Am. Midl. Nat.* 118, 358–365.
- Steuter, A. A., Jasch, B., Ihnen, J., and Tieszen, L. L. (1990). Woodland/grassland boundary changes in the middle Niobrara valley of Nebraska identified by $^{13}\text{C}/^{12}\text{C}$ values of soil organic matter. *Am. Midl. Nat.* 124, 301–308.
- Stokes, C. J. (1999). Woody plant dynamics in a south Texas savanna: pattern and process. Ph.D. Dissertation, Texas A&M University, College Station.

- Stroh, J. C. (1995). Landscape development and dynamics of a subtropical savanna parkland, 1941–1990. Ph.D. Dissertation, Texas A&M University, College Station.
- Thomas, D. B. and Pratt, D. J. (1967). Bush control studies in the drier areas of Kenya. IV. Effects of controlled burning on secondary thicket in upland *Acacia* woodland. *J. Appl. Ecol.* 4, 325–335.
- Tieszen, L. L. and Archer, S. R. (1990). Isotopic assessment of vegetation changes in grassland and woodland systems. In "Plant Biology of the Basin and Range" (C. B. Osmond, L. F. Pitelka, and G. M. Hidy, Eds.), pp. 293–321. Ecology Studies 80 Springer-Verlag, New York.
- Tieszen, L. L. and Pfau, M. W. (1995). Isotopic evidence for the replacement of prairie forest in the Loess Hills of eastern South Dakota. In "Proceedings, 14th North American Prairie Conference: Prairie Biodiversity" (D. C. Hartnett, Ed.), pp. 153–165.
- Tongway, D. and Ludwig, J. (1997). The nature of landscape dysfunction in rangelands. In "Landscape Ecology, Function and Management: Principles from Australia's Rangelands" (J. Ludwig, D. Tongway, D. Freudenberger, J. Noble, and K. Hodgkinson, Eds.), pp. 49–61. CSIRO Publishing, Melbourne.
- Tothill, J. C. and Mott, J. J. (Eds.). (1985). "Ecology and Management of the World's Savannas." Australian Academy of Science, Canberra, ACT.
- Trexler, M. C. and Meganck, R. (1993). Biotic carbon offset programs: sponsors of or impediment to economic development? *Climate Res.* 3, 29–136.
- Trollope, W. S. W. (1982). Ecological effects of fire in South African savannas. In "Ecology of Tropical Savannas" (B. J. Huntley and B. H. Walker, Eds.), pp. 292–306. Ecology Studies, No. 42, Springer-Verlag, Berlin.
- Turner, B. L. I., Clark, W. C., Kates, R. W., Richards, J. F., Mathews, J. T., and Meyers, W. B. (1990). "The Earth as Transformed by Human Action." Cambridge University Press, New York.
- Van Vegten, J. A. (1983). Thornbush invasion in a savanna ecosystem in eastern Botswana. *Vegetatio* 56, 3–7.
- Veble, T. T. and Lorenz, D. C. (1991). "The Colorado Front Range: A Century of Ecological Change." University of Utah Press, Salt Lake City.
- Vega, A. J. (1991). Simulating the hydrology and plant growth of south Texas rangelands. Ph.D. Dissertation, Texas A&M University, College Station.
- Virginia, R. A. (1986). Soil development under legume tree canopies. *Forest Ecol. Manage.* 16, 69–79.
- Virginia, R. A., Jenkins, M. B., and Jarrell, W. M. (1986). Depth of root symbiont occurrence in soil. *Biol. Fertil. Soils* 2, 127–130.
- Vitousek, P. M. (1994). Beyond global warming: Ecology and global change. *Ecology* 75, 1861–1876.
- Vivrette, N. J. and Muller, C. H. (1977). Mechanism of invasion and dominance of coastal grassland by *Mesembryanthemum crystallinum*. *Ecol. Monogr.* 47, 301–318.
- Walker, B. H. (Ed.). (1987). "Determinants of Tropical Savannas." IRL Press, Oxford, United Kingdom.
- Walker, B. H. (1993a). Rangeland ecology—understanding and managing change. *Ambio* 22, 80–87.
- Walker, B. H. (1993b). Stability in rangelands: ecology and economics. In "Proceedings, XVII International Grassland Congress" (M. J. Baker, J. R. Crush, and L. R. Humphreys, Eds.), pp. 1885–1890. New Zealand/Australia, Keeling & Monday, Palmerston North.
- Walker, B. H., Ludwig, D., Holling, C. S., and Peterman, R. M. (1981). Stability of semi-arid savanna grazing systems. *J. Ecol.* 69, 473–498.
- Walker, B. H. and Steffen, W. L. (1993). Rangelands and global change. *Rangel. J.* 15, 95–103.
- Wall, T. (1999). Western juniper encroachment into aspen communities in the northwest Great Basin. M.S. Thesis, Oregon State University, Corvallis.
- Wang, Y., Cerling, T. E., and Effland, W. R. (1993). Stable isotope ratios of soil carbonate and soil organic matter as indicators of forest invasion of prairie near Ames, Iowa. *Oecologia* 95, 365–369.
- Watts, S. E. (1993). Rooting patterns of co-occurring woody plants on contrasting soils in a subtropical savanna. M.S. thesis, Texas A&M University, College Station.
- Weltzin, J. F., Archer, S., and Heitschmidt, R. K. (1997). Small-mammal regulation of vegetation structure in a temperate savanna. *Ecology* 78, 751–763.
- Werner, P. A. (1991). "Savanna Ecology and Management: Australian Perspectives and Intercontinental Comparisons." Blackwell Science, London.
- West, N. E. (1986). Desertification or xerification? *Nature* 321, 562–563.
- West, N. E. (1988). Inter-mountain deserts, shrubsteppes and woodlands. In "North American Terrestrial Vegetation" (M. G. Barbour and W. D. Billings, Eds.), pp. 209–230. Cambridge University Press, New York.
- West, O. (1947). Thorn bush encroachment in relation to the management of veld grazing. *Rhodesia Agri. J.* 44, 488–497.
- Williams, K., Hobbs, R. J., and Hamburg, S. P. (1987). Invasion of an annual grassland in Northern California by *Baccharis pulularis* sp. *consanguinea*. *Oecologia* 72, 461–465.
- Wondzell, S. and Ludwig, J. A. (1995). Community dynamics of desert grasslands: influences of climate, landforms and soils. *J. Veg. Sci.* 6, 377–390.
- York, J. C. and Dick-Peddie, W. A. (1969). Vegetation changes in southern New Mexico during the past one-hundred years. In "Arid Lands in Perspective" (W. G. McGinnis and B. J. Goldman, Eds.), pp. 155–166. University of Arizona Press, Tucson.
- Yorks, T. P., West, N. E., and Capels, K. M. (1992). Vegetation differences in desert shrublands of western Utah's Pine Valley between 1933 and 1989. *J. Range Manage.* 45, 569–578.
- Young, J. A., Eckert, R. E., and Evans, R. A. (1979). Historical perspectives regarding the sagebrush ecosystem. In "The Sagebrush Ecosystem: A Symposium." pp. 1–13. College of Natural Resources, Utah State University, Logan.
- Young, J. A. and Evans, R. A. (1981). Demography and fire history of a western juniper stand. *J. Range Manage.* 34, 501–506.
- Young, M. D. and Solbrig, O. T. (1993). "The World's Savannas: Economic Driving Forces, Ecological Constraints and Policy Options for Sustainable Land Use." Parthenon Publishing Group, Carnforth, United Kingdom.
- Zak, D. R., Tilman, D., Parmenter, R. R., Rice, C. W., Fisher, F. M., Vose, J., Milchunas, D., and Martin, C. W. (1994). Plant production and soil microorganisms in late-successional ecosystems: a continental-scale study. *Ecology* 75, 2333–2347.
- Zimmerman, G. T. and Neunshwander, L. F. (1984). Livestock grazing influences on community structure, fire intensity and fire frequency within the douglas-fir/ninebark habitat type. *J. Range Manage.* 37, 104–110.
- Zitzer, S. F., Archer, S. R., and Boutton, T. W. (1996). Spatial variability in the potential for symbiotic N_2 fixation by woody plants in a subtropical savanna ecosystem. *J. Appl. Ecol.* 33, 1125–1136.




CHAPTER 18, FIGURE 1 Upper panels: simplified scheme of the natural global cycles of carbon (left) and atmospheric molecular oxygen (right). Ph, photosynthesis; Res, respiration; MB, marine biota; DIC, dissolved inorganic carbon (H_2CO_3 , HCO_3^- , CO_3^{2-}). Pool sizes are not shown to scale. Lower panels: corresponding schemes of the cycles of the oxygen isotopes in CO_2 (lower left panel) and in O_2 (lower right panel). Dark blue arrows indicate links to oxygen isotopes in the hydrological cycle. Red whiskers on arrows indicate exchanges during which fractionation processes occur.



CHAPTER 18, FIGURE 2 Globally averaged O_2/N_2 ratio (vertical axis) versus globally averaged CO_2 mixing ratio (horizontal axis). The annually averaged observations have been determined from the station records as described in the text. The black arrow shows the observed trend for 1990–1997 determined by the fitting procedure as described in the text. The red arrow depicts the expected change due to the fossil emissions during 1990–1997. The effects of the ocean and the land biosphere is shown with the blue and the green arrow, whereby their slopes are determined by their respective O_2 versus CO_2 contributions (see inset). The purple vertical arrow reflects an estimate of the oceanic O_2 outgassing induced by ocean warming.



CHAPTER 9, FIGURE 12 LANDSAT image (1978) of the Canada (Saskatchewan/Alberta)–U.S. (Montana) border in the vicinity of the Milk River (from Knight, 1991). Subsequent to the drought and Dust Bowl of the 1930s, farmlands in Canada were repossessed by provincial or federal governments, withdrawn from cultivation, and underwent secondary succession. Intensive agriculture was maintained in the United States via elaborate farm subsidy programs. Striking contrasts in regional land cover were thus a direct result of changes in government policy.



CHAPTER 16, FIGURE 3 Potential mean soil carbon turnover rates extrapolated to the global scale using the temperature and soil texture relationships from the Century model (Schimel *et al.*, 1994).