



## Scouting Sugarcane Aphids

Robert Bowling, Michael Brewer, Allen Knutson, Mo Way, Pat Porter, Ed Bynum, Charles Allen, Raul Villanueva\*

\*All of the Texas A&M AgriLife Extension Service, except Brewer and Way, of Texas A&M AgriLife Research.

Immature Sugarcane Aphid



Winged Adults



Timing effective treatment to control sugarcane aphids (SCA) in sorghum depends on the size of the SCA population. To estimate the number of SCA in a field, follow these steps for scouting the field and use the **Sampling Protocol** (below) and the **Quick Aphid Checker** (on back) to make treatment decisions.

### First Detection: Is the Field at Risk?

- Once a week, walk 25 feet into the field and examine plants along 50 feet of row.
- If honeydew is present, look for SCA on the underside of a leaf above the honeydew.
- Inspect the underside of leaves from the upper and lower canopy from 15–20 plants per location.
- Sample each side of the field as well as sites near Johnsongrass and tall mutant plants.
- Check at least 4 locations per field for a total of 60–80 plants.

If no SCA are present, or only a few wingless/winged aphids are on upper leaves, continue once-a-week scouting.

If SCA are found on lower or mid-canopy leaves, begin twice-a-week scouting. Use the Sampling Protocol and the Quick Aphid Checker to determine if aphid densities exceed the economic threshold.

### Sampling Protocol: Making Treatment Decisions

Examine the underside of one completely green leaf from the lower canopy and the uppermost leaf (or the leaf below the flag leaf at boot to heading) and estimate the number of SCA per leaf, using the Quick Aphid Checker. Examine 2 leaves from each of 5 random plants per location. Repeat at 4 locations, for a total of 40 leaves. Use the Quick Aphid Checker to calculate the mean number of aphids per leaf.

- If the field average SCA infestation is **50–125 aphids or more per leaf**, apply an insecticide within 4 days and evaluate control after 3–4 days. Consider treatment at 50 aphids per leaf if limited to once-a-week scouting.
- If the SCA infestation is less than the threshold level, continue scouting twice a week.



# Quick Aphid Checker

Estimate the number of sugarcane aphids (SCA) per leaf to help time foliar insecticides for SCA control on sorghum. Each photo represents an estimate from the table. For example, photo A shows about 12 aphids.

Estimate the Number of Aphids per Leaf		
Photo	Range	Estimate
A	1-25	12
B	26-50	38
C	51-100	75
D	101-500	300
E	501-1000	750
F	>1000	1500

Field Average =  $\frac{\text{Total of All Estimates}}{\text{Total \# of Leaves Examined}}$

Learn more about sugarcane aphids at <http://txscan.blogspot.com>

*Photos courtesy of Travis Ahrens, Mike Brewer, and Pat Porter.  
Funding provided by the Texas Grain Sorghum Producers Board and the USDA NIFA Southern IPM Center and Crop Protection and Pest Management Program.*

Educational programs of the Texas A&M AgriLife Extension Service are open to all people without regard to race, color, sex, religion, national origin, age, disability, genetic information, or veteran status.

1000 copies – New

